



Editor's choice
Scan to access more
free content

Which determinants predict tibiofemoral and patellofemoral osteoarthritis after anterior cruciate ligament injury? A systematic review

Belle L van Meer,¹ Duncan E Meuffels,¹ Wilbert A van Eijsden,¹ Jan A N Verhaar,¹ Sita M A Bierma-Zeinstra,^{1,2} Max Reijnen¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2013-093258>).

¹Department of Orthopaedic Surgery, Erasmus MC, University Medical Centre Rotterdam, The Netherlands
²Department of General Practice, Erasmus MC, University Medical Centre Rotterdam, The Netherlands

Correspondence to

Belle L van Meer, Department of Orthopaedic Surgery, Erasmus MC, University Medical Centre Rotterdam, PO Box 2040, 3000 CA Rotterdam, The Netherlands; b.vanmeer@erasmusmc.nl

Accepted 2 March 2015
Published Online First
30 March 2015

ABSTRACT

Background Anterior cruciate ligament (ACL) injury is an important risk factor for development of knee osteoarthritis (OA). To identify those ACL injured patients at increased risk for knee OA, it is necessary to understand risk factors for OA.

Aim To summarise the evidence for determinants of (1) tibiofemoral OA and (2) patellofemoral OA in ACL injured patients.

Methods MEDLINE, EMBASE, Web of Science and CINAHL databases were searched up to 20 December 2013. Additionally, reference lists of eligible studies were manually and independently screened by two reviewers. 2348 studies were assessed for the following main inclusion criteria: ≥ 20 patients; ACL injured patients treated operatively or non-operatively; reporting OA as outcome; description of relationship between OA outcome and determinants; and a follow-up period ≥ 2 years. Two reviewers extracted the data, assessed the risk of bias and performed a best-evidence synthesis.

Results Sixty-four publications were included and assessed for quality. Two studies were classified as low risk of bias. Medial meniscal injury/menisectomy showed moderate evidence for influencing OA development (tibiofemoral OA and compartment unspecified). Lateral meniscal injury/menisectomy showed moderate evidence for no relationship (compartment unspecified), as did time between injury and reconstruction (tibiofemoral and patellofemoral OA).

Conclusions Medial meniscal injury/menisectomy after ACL rupture increased the risk of OA development. In contrast, it seems that lateral meniscal injury/menisectomy has no relationship with OA development. Our results suggest that time between injury and reconstruction does not influence patellofemoral and tibiofemoral OA development. Many determinants showed conflicting and limited evidence and no determinant showed strong evidence.

symptomatic relief, osteotomy, unicompartmental arthroplasty and, for end-stage disease, total knee arthroplasty. Early intervention is critical because patients with post-traumatic OA are typically young and it is important to postpone total knee arthroplasty.⁴

Numerous studies have evaluated the long-term consequences of ACL rupture. These studies are heterogeneous with regard to methodology, including treatments used, inclusion of additional intra-articular injuries, reported OA outcomes and descriptions of determinants (potential risk factors). Three previous systematic reviews of development of OA after ACL rupture were limited either because they considered OA only in the tibiofemoral compartment or because they focused on one type of treatment (ACL reconstruction). Oiestad *et al*⁵ conducted a systematic review of the prevalence of OA in the tibiofemoral joint occurring more than 10 years after ACL injury. They included studies that used ACL reconstruction techniques, which are no longer used (eg, Leeds-Keio polyester ligament surgery or suturing of the ACL). Therefore, we did not include these techniques in this systematic review. To better evaluate newer and current techniques and rehabilitation methods, we included only studies which reported results based on current ACL reconstruction procedures. Magnussen and Spindler⁶ reviewed patient factors affecting clinical and radiographic outcomes after ACL reconstruction in prospective studies with a 5-year minimum follow-up. Prospective study design was an inclusion criterion, so they missed the results of all retrospective studies. Claes *et al*⁷ reviewed the literature on long-term radiographic outcome after autologous ACL reconstruction; studies with a mean follow-up of less than 10 years were excluded. They investigated only one predictor, namely the relationship between meniscal status and OA development in the reconstructed knee. Currently, there is no consensus about operative or non-operative treatment for preventing OA, and degenerative changes can develop in all knee compartments.

Culvenor *et al*⁸ showed in their narrative literature review that patellofemoral OA after ACL reconstruction occurs as frequently as tibiofemoral OA. Different mechanisms, such as inflammation, concomitant injuries to the patellofemoral articular cartilage, meniscal injury, graft choice and changes of knee biomechanics, may play a role in the development of patellofemoral OA.⁸

The previous published reviews presented a part of the general question: Which determinants influence the development of degenerative changes after

INTRODUCTION

Anterior cruciate ligament (ACL) rupture is a common sports-related injury, with an annual incidence of approximately 5/10 000 persons in the general population.¹ Osteoarthritis (OA) is a well-known, long-term complication of ACL rupture, with a prevalence of 10–90% at 10–20 years post-injury.^{2–3} It is important to identify the risk factors contributing to OA in patients with ACL rupture, because some risk factors may be modifiable as to prevent onset or early-stage progression of OA. At present, the only treatment options for OA are



CrossMark

To cite: van Meer BL, Meuffels DE, van Eijsden WA, *et al*. *Br J Sports Med* 2015;**49**: 975–983.

an ACL rupture? This systematic review will fill the gaps of the previous reviews and supplement recent published literature on both tibiofemoral and patellofemoral OA. We systematically reviewed the evidence for determinants of both (1) tibiofemoral OA and (2) patellofemoral OA in patients with an ACL injury treated operatively or non-operatively.

METHODS

The reporting in this systematic review was conducted according to the PRISMA statement.⁹

Data sources and searches

MEDLINE, EMBASE, Web of Science and CINAHL medical literature databases were searched up to 20 December 2013. Search terms included anterior cruciate ligament, synonyms for injury and synonyms for osteoarthritis. The full electronic search strategy for the MEDLINE database is presented in table 1. Similar search strategies were used in EMBASE, Web of Science and CINAHL. Additionally, the reference lists of all eligible studies were manually screened.

Study selection

Two reviewers (BLvM and MR) assessed the studies for the following inclusion criteria:

- ▶ The following study designs with at least 20 patients: randomised controlled trial, prospective follow-up study, matched case-control study and retrospective study;
- ▶ Subjects had to have an ACL injury consisting of:
 - Patients treated non-operatively or
 - Patients treated operatively; use of an arthroscopic or miniarthrotomy technique and use of bone-patellar tendon-bone, hamstring tendon or allografts;
- ▶ Written in English, German, Dutch, Spanish, French, Swedish, Danish or Norwegian;
- ▶ Full text available;
- ▶ Measured one of the following OA outcomes:
 - Clinical OA: according to a clinician, self-reported or American College of Rheumatology (ACR) criteria¹⁰; osteotomy, unilateral knee arthroplasty or total knee arthroplasty (indirect measures for clinical knee OA);
 - Radiographic OA;
 - OA findings on MRI;
 - OA findings during arthroscopy;
- ▶ The relationship between outcome and determinant, defined as potential risk factors, must have been described or data must be available to calculate the relationship;
- ▶ Determinant studied in ≥ 2 studies;
- ▶ Determinant must be measured prior to the OA outcome;
- ▶ Follow-up period of at least 2 years.

Animal studies and reviews were excluded. Disagreements on inclusions were resolved by discussion and, if necessary, a final decision was made by a third reviewer (JANV).

Data extraction and risk of bias assessment

Two reviewers (WAvE and BLvM) extracted the study characteristics, follow-up times, determinants, outcomes and the relationship between outcome and determinant.

The determinants were grouped into *patient characteristics* (age, body mass index (BMI), sex), *physical examination, activity level* and *intra-articular-related factors*. The determinant that was named laxity consisted of results of a pivot shift test, Lachman test, KT 1000 arthrometer or description of 'laxity'. The location of injury of the intra-articular determinants: chondral injury and meniscal injury/meniscectomy were presented when reported as such in the studies. For determining the influence of *tunnel placement* on OA development, we used the assessment of tunnel position when a study evaluated both femoral and tibial tunnel positions and graft inclination. If studies had the same population and determinant, but different follow-up times, we presented the results of the study with the longest follow-up time. When a determinant was measured in various ways and had different relationships with OA outcome in one study, all results were presented. For the analyses of the relationship between determinants and OA outcome, the distinction between patellofemoral and tibiofemoral OA was made. If the studies did not report a specific compartment for the OA outcome or if the studies reported the OA outcome for all compartments, then the study was classified as OA outcome in which the compartment was unspecified. Since the included studies presented the relationship between determinant and OA outcome in various ways, we reported the presence of a 'positive significant relationship' or 'negative significant relationship' or 'no significant relationship'. For presentation of the results, we distinguished the studies into two groups: (1) studies with inclusion of non-operatively treated patients and (2) studies with inclusion of both operatively and non-operatively or solely operatively treated patients.

We evaluated the selected studies on 12 aspects using modified questions of existing risk of bias assessment tools.^{11–13} Our assessment tool contained questions about the aim of the study, description of inclusion and exclusion criteria, collection of data, validity and reliability of OA outcome measures, independent measure of determinants, valid and reliable measurement of determinants, follow-up period, loss to follow-up, and use of adequate statistical analyses. Four reviewers independently assessed the quality of the included studies. Disagreements were resolved by discussion. Studies were classified as low risk of bias when they scored 'adequate' on all the following topics: the authors reported inclusion of consecutive patients; there was unbiased assessment of the study outcome and determinants; the determinant measures were used accurately (valid and reliable); if there was a loss to follow-up of less than 20% and there was a description of the reasons, and if there was correction for confounding. The assessment tool used is given in online supplementary appendix table S1.

Data synthesis and analysis

Since the studies were considered clinically heterogeneous with regard to the outcome measures and determinants studied, it was not possible to pool the data for statistical analysis, and therefore we performed 'a best-evidence synthesis'.^{14 15} With the use of the system developed by van Tulder *et al*,¹⁶ the following ranking of levels of evidence was formulated: (1) Strong evidence is provided by two or more studies with low risk of

Table 1 Search strategy for MEDLINE

```
(anterior cruciate*[tw] OR acl[tw])
AND
(rupture*[tw] OR tear*[tw] OR torn*[tw] OR lacerat*[tw] OR defici*[tw] OR
injur*[tw] OR lesion*[tw] OR disrupt*[tw] OR trauma*[tw] OR reconstruct*[tw]
OR repair*[tw])
AND
(osteoarthritis*[tw] OR osteo-arthritis*[tw] OR osteoarthro*[tw] OR osteo-arthro*
[tw] OR arthrosis[tw] OR arthroses[tw] OR arthrot*[tw] OR gonarthro*[tw] OR
degen*[tw])
NOT
(animals[mesh] NOT humans[mesh])
```



bias and by generally consistent findings in all studies ($\geq 75\%$ of the studies reported consistent findings). (2) Moderate evidence is provided by one low risk of bias study and two or more high risk of bias studies and by generally consistent findings in all studies ($\geq 75\%$). (3) Limited evidence is provided by one or more high risk of bias studies or one low risk of bias study and by generally consistent findings ($\geq 75\%$). (4) Conflicting evidence is provided by conflicting findings ($< 75\%$ of the studies reported consistent findings). (5) No evidence is provided when no studies could be found.^{16 17}

RESULTS

Identification and selection of the literature

The search resulted in 2348 studies, for which all abstracts were reviewed. After screening of the abstracts, 157 were identified as possibly relevant, and full texts were retrieved. After review of the full texts, 56 met all the inclusion criteria (figure 1). There were no disagreements on inclusions. The references of all 56 studies were reviewed and 8 additional studies meeting the inclusion criteria were identified. Thus, 64 studies in total were included in this systematic review.

Figure 1 Study selection (ACL, anterior cruciate ligament; OA, osteoarthritis).



Description of the included studies

The characteristics of the included studies are presented in online supplementary appendix table S2. The studies had the following designs: randomised controlled trial (n=12),^{18–29} prospective follow-up study (n=22),^{30–51} matched case-control study (n=2)^{52–53} and retrospective study (n=28).^{54–81} The number of patients available for follow-up measurement in the studies ranged between 30 and 780. In 62 studies, the OA outcome was determined with radiographs, and in 2 studies by MRI assessment.^{28–47} Only two studies^{43–70} reported radiological OA and clinical OA as outcomes. Therefore, the findings of this systematic review address the influence of radiological OA. In 47 studies (4956 patients), the treatment strategy was ACL reconstruction, in 4 studies^{22–64–71–76} (273 patients) non-operative treatment, and in 13 studies^{19–30–31–40–41–47–53–65–70–72–77–79–80} (1169 patients) both reconstruction and non-operative treatment. The mean follow-up time varied between 3.9 and 20 years.

Risk of bias assessment

Two studies^{35–56} were classified as ‘low-risk of bias’. An overview of the quality assessment score of the included studies is presented in supplementary appendix table S3. The main aim of the two low risk of bias studies was to investigate risk factors for development of knee OA after ACL reconstruction. In these

studies, the number of patients used for analyses was >50; Ahn *et al* had a sample size of more than 100 patients (n=117). Janssen *et al* used only hamstring tendon grafts, and Ahn *et al* bone-patellar tendon-bone grafts.

Influence of determinants in non-operatively treated patients

Four studies^{22–64–71–76} included solely non-operatively treated patients. Limited evidence was found for a positive relationship between *meniscectomy* and development of knee OA in chronic ACL-deficient knees. Determinants *age*, *BMI* and *sex* were excluded because they were studied in only one study. The influence of *laxity* on OA development could not be presented because it was measured concurrently with the OA outcome.

Influence of determinants in both operatively and non-operatively or solely operatively treated patients

Patient characteristics

Conflicting evidence was found for the influence of *age* on OA outcome in all compartments (tables 2–4). For the influence of *BMI* on OA outcome in the tibiofemoral compartment and compartment unspecified, conflicting evidence was found after ACL rupture. Limited evidence for no relationship was found for OA development in the patellofemoral compartment after ACL

Table 2 Influence of determinants on tibiofemoral radiological OA outcome in operatively and operatively/non-operatively treated cohorts

Group	Determinant	Number of studies	Significant relationship LR/HR: n studies	No significant relationship LR/HR: n studies	Best-evidence synthesis	Comments
Patient characteristics	Older age	6	Positive relationship: LR: 2 ^{43–66}	LR: 1 ⁵⁶ HR: 3 ^{37–40–60}	Conflicting evidence	
	Higher BMI	4	Positive relationship: LR: 1 ^{56*} HR: 1 ⁶⁶	HR: 2 ^{40–43}	Conflicting evidence	*Ahn <i>et al</i> : ⁵⁶ lateral OA
	Male sex	3	Positive relationship: HR: 1 ⁴³	HR: 2 ^{60–66}	Conflicting evidence	
Intra-articular-related factors	Additional injury	2	Positive relationship: HR: 1 ⁴⁴	HR: 1 ⁶³	Conflicting evidence	
	Chondral injury	3	Positive relationship: HR: 2 ^{37–60}	HR: 1 ⁶⁶	Conflicting evidence	
	Meniscal injury/meniscectomy	11	Positive relationship: Medial meniscus injury/meniscectomy: LR: 1 ⁵⁶ HR: 2 ^{20–57} Lateral meniscus injury/meniscectomy: HR: 1 ⁵⁷ Location not reported: HR: 6 ^{37–40–60–69–80–81}	Medial meniscus injury/meniscectomy: None Lateral meniscus injury/meniscectomy: LR: 1 ³⁶ HR: 1 ²⁰ Location not reported: HR: 2 ^{48–66}	Medial meniscus injury/meniscectomy: Moderate evidence positive relationship Lateral meniscus injury/meniscectomy: Conflicting evidence Location not reported: Limited evidence positive relationship	
	Longer time between injury and reconstruction	6	Positive relationship HR: 1 ⁶¹	LR: 1 ⁵⁶ HR: 4 ^{37–43–60–66}	Moderate evidence for no relationship	
	ACL reconstruction (vs non-operative treatment)	7	Positive relationship: HR: 2 ^{31–40} Negative relationship: HR: 1 ⁴⁷	HR: 4 ^{19–53–70–80}	Conflicting evidence	
	Graft type BPTB (vs HT)	8	Positive relationship HR: 4 ^{37–38–43–52}	HR: 6 ^{**18–19–21–38–52–69}	Conflicting evidence	*Leys <i>et al</i> : ³⁸ medial OA; Mascarenhas <i>et al</i> : ⁵² lateral OA **Leys <i>et al</i> : ³⁸ lateral OA; Mascarenhas <i>et al</i> : ⁵² medial OA

*Refers to the comment in the last column in the same row.

ACL, anterior cruciate ligament; BMI, body mass index; BPTB, bone-patellar tendon-bone; HR, high risk of bias studies; HT, hamstring tendon; LR, low risk of bias studies; OA, osteoarthritis.

Table 3 Influence of determinants on patellofemoral radiological OA outcome in operatively and operatively/non-operatively treated cohorts

Group	Determinant	Number of studies	Significant relationship LR/HR: n studies	No significant relationship LR/HR: n studies	Best-evidence synthesis	Comments
Patient characteristics	Older age	3	Positive relationship: HR: 2 ^{37 42}	LR: 1 ⁵⁶	Conflicting evidence	
	Higher BMI	2		LR: 1 ⁵⁶ HR: 1 ⁴²	Limited evidence for no relationship	
Intra-articular	Meniscal injury/meniscectomy	3	Positive relationship: HR: 1 ⁴¹	LR: 1 ⁵⁶ HR: 1 ³⁷	Conflicting evidence	
	Longer time between injury and reconstruction	3		LR: 1 ⁵⁶ HR: 2 ^{37 42}	Moderate evidence for no relationship	
	ACL reconstruction (vs non-operative treatment)	5	Positive relationship: HR: 2 ^{31 41}	HR: 3 ^{19 47 70}	Conflicting evidence	
	Graft type BPTB (vs HT)	6	Positive relationship: HR: 1 ¹⁹	HR: 5 ^{18 37 38 52 69}	Limited evidence for no relationship	
	Tunnel placement	2		LR: 1 ⁵⁶ HR: 1 ⁶²	Limited evidence for no relationship	

ACL, anterior cruciate ligament; BMI, body mass index; BPTB, bone-patellar tendon-bone; HR, high risk of bias studies; HT, hamstring tendon; LR, low risk of bias studies; OA, osteoarthritis.

rupture. Nine studies evaluated the relationship between *sex* and OA development after ACL rupture. For development of tibiofemoral OA, three high risk of bias studies^{43 60 66} showed conflicting evidence. Moderate evidence was found for no relationship between male sex and OA development in compartment unspecified.^{25 35 67 68 75 79}

Physical examination

One low risk of bias³⁵ and two high risk of bias^{34 45} studies showed no relationship between *laxity* and development of OA in compartment unspecified (table 4). Thus, there is moderate evidence for no relationship between laxity and OA development.^{34 35 45} Moderate evidence was also found for no relationship between range of motion and OA development in compartment unspecified.^{34 35 45 50} Performance of a single-legged hop test was evaluated in three studies^{34 35 45} and showed conflicting evidence.

Activity level

One low risk of bias study³⁵ and one high risk⁶⁸ of bias study found no significant relationship between activity level before reconstruction and OA development (compartment unspecified; table 4).

Intra-articular-related factors

Two high risk of bias studies^{44 63} investigating *additional injuries in general* showed conflicting evidence (tables 2–4).

One high risk of bias study⁶⁸ evaluated patellar, medial and lateral chondral injury after ACL rupture and their influence on OA development in compartment unspecified. Medial and patellar chondral injury showed a positive significant relationship with development of knee OA and lateral chondral injury showed no relationship. There were ten other studies,^{23 25 33 35 37 49 50 60 66 73} of which one low risk of bias study³⁵ showed conflicting evidence if the location of the chondral injury was not reported.

In nine studies,^{20 35 49 55–57 67 68 75} of which two were low risk of bias studies, a distinction between medial and lateral *meniscus injury/meniscectomy* was made. We found moderate evidence for a positive relationship between medial meniscus injury/meniscectomy and development of OA (tibiofemoral and unspecified) in patients with an ACL rupture. Conflicting evidence was found for influence of lateral meniscus injury/

meniscectomy on tibiofemoral OA development and moderate evidence for no significant relationship on OA development in compartment unspecified. Twenty-one high risk of bias studies did not report the location of the meniscus injury; these studies showed limited evidence for positive relationship with development of tibiofemoral OA and conflicting evidence if the compartment of OA development was unspecified. The studies did not report the extent of meniscectomy. Results of meniscus injury/meniscectomy showed conflicting evidence for a relationship with patellofemoral OA development. One low risk of bias study⁵⁶ and one high risk³⁷ of bias study reported no significant relationship, and in one high risk of bias study⁴¹ meniscus injury/meniscectomy was related to patellofemoral OA development.

In seven studies,^{37 42 43 56 60 61 66} one of them low risk of bias study, moderate evidence for no relationship was found for the influence of *time between injury and reconstruction* on development of tibiofemoral and patellofemoral OA. Seven studies did not specify the compartment of OA outcome and these studies showed conflicting evidence.^{25 35 36 68 74 78 79}

In 13 studies investigating *ACL reconstruction* versus *non-operative treatment*, conflicting evidence was found with patellofemoral OA,^{19 31 41 47 70} tibiofemoral OA^{19 31 40 53 70 80} and if no specific compartment^{30 65 72 77 79} was reported.

Fourteen studies reported outcomes on the relationship between *bone-patellar tendon-bone graft* versus *hamstring tendon graft* and development of tibiofemoral OA or OA in compartment unspecified. The studies gave conflicting findings. Mascarenhas *et al*⁵² and Leys *et al*³⁸ reported opposite results for the development of medial and lateral tibiofemoral OA; Mascarenhas *et al* found a positive relationship between bone-patellar tendon-bone graft and development of lateral tibiofemoral OA, whereas Leys *et al* found a positive relationship between bone-patellar tendon-bone graft and development of medial tibiofemoral OA. In six studies,^{18 19 37 38 52 69} the influence of graft type on patellofemoral OA was studied: limited evidence was found for no relationship. Conflicting evidence in two high risk of bias studies^{26 68} was found for the influence of *allograft* on OA development in compartment unspecified.

One low risk of bias study and five high risk of bias studies reported on the influence of *tunnel placement* of the ACL reconstruction and OA development. Two studies showed no

Table 4 Influence of determinants on radiological OA outcome compartment unspecified in operatively and operatively/non-operatively treated cohorts

Group	Determinant	Number of studies	Significant relationship LR/HR: n studies	No significant relationship LR/HR: n studies	Best-evidence synthesis	Comments
Patient characteristics	Older age	9	Positive relationship: HR: 3 ^{50 65 78}	LR: 1 ³⁵ HR: 5 ^{25 67 68 75 79}	Conflicting evidence	
	Higher BMI	5	Positive relationship: HR: 2 ^{65 68}	LR: 1 ³⁵ HR: 2 ^{67 79}	Conflicting evidence	
	Male sex	6	Positive relationship: HR: 1 ⁶⁸	LR: 1 ³⁵ HR: 4 ^{25 67 75 79}	Moderate evidence for no relationship	
Physical examination	Laxity	3		LR: 1 ³⁵ HR: 2 ^{34 45}	Moderate evidence for no relationship	
	Range of motion loss	4	Positive relationship: HR: 1 ⁵⁰	LR: 1 ³⁵ HR: 2 ^{34 45}	Moderate evidence for no relationship	
	Performance single-legged hop test	3	Negative relationship: HR: 1 ⁴⁵	LR: 1 ³⁵ HR: 1 ³⁴	Conflicting evidence	
Activity	Activity level before reconstruction	2		LR: 1 ³⁵ HR: 1 ⁶⁸	Limited evidence for no relationship	
Intra-articular	Chondral injury	8	<i>Medial chondral injury:</i> Positive relationship: HR: 1 ⁶⁸ <i>Lateral chondral injury:</i> None <i>Patellar chondral injury:</i> Positive relationship: HR: 1 ⁶⁸ <i>Location not reported</i> Positive relationship: LR: 1 ³⁵ HR: 3 ^{33 50 73}	<i>Medial chondral injury:</i> None <i>Lateral chondral injury:</i> HR: 1 ⁶⁸ <i>Patellar chondral injury:</i> None <i>Location not reported</i> HR: 3 ^{23 25 49}	<i>Medial chondral injury:</i> Limited evidence for positive relationship <i>Lateral chondral injury:</i> Limited evidence for no relationship <i>Patellar chondral injury:</i> Limited evidence for positive relationship <i>Location not reported</i> Conflicting evidence	
	Meniscal injury/meniscectomy	19	Positive relationship: Medial meniscus injury/meniscectomy: LR: 1 ³⁵ HR: 5 (49, 55, 67, 68: concurrent*, 75) Lateral meniscus injury/meniscectomy: None Both meniscectomy: 1 ⁴⁹ <i>Location not reported:</i> HR: 9 ^{36 39 50 51 54 70 73 74 78}	Medial meniscus injury/meniscectomy: HR: 1 (68: prior*) Lateral meniscus injury/meniscectomy: LR: 1 ³⁵ HR: 4 (49, 67, 68: prior and concurrent, 75) Both meniscectomy: None <i>Location not reported:</i> HR: 4 ^{23 32 59 79}	Medial meniscus injury/meniscectomy: Moderate evidence positive relationship Lateral meniscus injury/meniscectomy: Moderate evidence no relationship Both meniscectomy: Limited evidence for positive relationship <i>Location not reported:</i> Conflicting evidence	*Li et al. ⁶⁸ concurrent: meniscectomy concurrent with ACL reconstruction; prior: meniscectomy prior to ACL reconstruction
	Longer time between injury and reconstruction	7	Positive relationship HR: 4 ^{36 68 74 78}	LR: 1 ³⁵ HR: 2 ^{25 79}	Conflicting evidence	
	ACL reconstruction (vs non-operative treatment)	5	Positive relationship: HR: 1 ⁶⁵ Negative relationship: HR: 2 ^{72 77}	HR: 2 ^{30 79}	Conflicting evidence	
	Graft type BPTB (vs HT)	7	Positive relationship: HR: 5 ^{24 29 38 45 68}	HR: 2 ^{23 28}	Conflicting evidence	
	Graft type allograft (vs autograft)	2	Positive relationship: HR: 1 ⁶⁸	HR: 1 ²⁶	Conflicting evidence	
	Tunnel placement	4	Positive relationship: HR: 1 ⁶⁸	HR: 3 ^{38 46 58}	Limited evidence for no relationship	
	Single bundle (vs double bundle)	2		HR: 2 ^{25 27}	Limited evidence for no relationship	

*Refers to the comment in the last column in the same row.

ACL, anterior cruciate ligament; BMI, body mass index; BPTB, bone-patellar tendon-bone; HT, hamstring tendon; HR, high risk of bias studies; LR, low risk of bias studies; OA, osteoarthritis.

significant relationship between tunnel placement and patellofemoral OA development.^{56 62} Four high risk of bias studies^{38 46 58 68} evaluated the influence on development of OA in compartment unspecified; three studies^{38 46 58} found no significant relationship, resulting in limited evidence for no relationship.

Two studies with high risk of bias reported on the influence of double-bundle and single-bundle ACL reconstruction and OA development in compartment unspecified.^{25 27} These studies showed limited evidence for no relationship with development of OA.

DISCUSSION

We summarised the available evidence concerning which determinants influence the risk of OA after ACL rupture. Sixty-four studies were included, but 62 were classified as high risk of bias.

Key clinically relevant findings

There was moderate evidence for:

- ▶ Medial meniscal injury/meniscectomy influencing OA development (tibiofemoral OA and compartment unspecified).
- ▶ No relationship with time between injury and reconstruction and OA development in patellofemoral and tibiofemoral compartments.
- ▶ No relationship between OA development in unspecified compartments and the following determinants was found: sex, laxity, range of motion and lateral meniscal injury/meniscectomy.

There was limited evidence for influencing OA development by the following determinants:

- ▶ Medial and patellar chondral injury (compartment unspecified).
- ▶ Meniscal injury/meniscectomy if the location was not reported (tibiofemoral OA).
- ▶ Meniscectomy of both menisci (compartment unspecified).
- ▶ Meniscectomy in non-operatively treated patients.

The following determinants showed limited evidence for no relationship with OA development:

- ▶ BMI (patellofemoral OA).
- ▶ Graft type (patellofemoral OA).
- ▶ Activity level pre-reconstruction (compartment unspecified).
- ▶ Lateral chondral injury (compartment unspecified).
- ▶ Tunnel placement (patellofemoral OA and compartment unspecified).
- ▶ Single-bundle versus double-bundle ACL reconstruction technique (compartment unspecified).

Outcome measure—OA

Notably, most studies reported only radiological OA. Only two studies^{43 70} reported both radiological OA and clinical OA as outcomes for evaluating the influence of determinants. Thus, the findings of this systematic review address the influence on radiological OA but not on clinical OA. We were also interested in determinants that influence early degenerative changes; however, the majority of the included studies reported mid-term or long-term follow-up. A mean follow-up time ≤ 5 years was reported in only eight studies.

The role of the meniscus: keep or cut?

Many studies evaluated the influence of the meniscus on the development of OA. The majority of studies did not report the location of the tear, the extent of meniscectomy, and in which compartment the OA was developing. We had no information

about the influence of the time of the meniscus injury, also a possible confounder.

Although more extended, our results are in line with the findings of the previous reviews concerning meniscal injury and meniscectomy as risk factors for tibiofemoral OA development. However, these previous reviews did not distinguish between medial and lateral meniscal injuries/meniscectomies.

Our review provides important data that medial meniscal injury/meniscectomy showed a relationship with the development of OA, but lateral meniscal injury/meniscectomy did not. Anatomically, the medial meniscus is more rigid with less anterior posterior mobility than the more mobile lateral meniscus; this could have an effect on the secondary OA changes of the affected compartment.⁸²

These findings contradict the results of a systematic review concerning clinical outcome and risk of OA development in patients undergoing meniscectomy. In that review, Salata *et al*⁸³ found four studies with a higher rate of OA in the lateral meniscectomy group, two studies reporting no significant difference, and one study in which medial meniscectomy was more related with OA. These results were not included in our systematic review because the meniscus studies did not meet the inclusion criteria. Moreover, most studies did not report the location (medial or lateral compartment) of the meniscal resection, making it difficult to discern the specific influence of medial/lateral meniscectomy.

A possible explanation for conflicting evidence for development of OA (compartment unspecified) and limited evidence for a positive relationship with development of tibiofemoral OA is the heterogeneity of the location of meniscectomy. Also, the included studies did not report the extent of meniscectomy, except the study of Fink *et al*,³⁰ which found in patients treated non-operatively for their ACL rupture a significant correlation between the degree of OA and the amount of meniscal resection that was performed at the time of the initial arthroscopy. For the ACL reconstructed group, there was no significant correlation.

A focus on patellofemoral OA

Patellofemoral OA is gaining consideration as an important clinical entity.⁸⁴ Regarding OA of the patellofemoral joint, two studies^{37 56} found no relationship with meniscal injury/meniscectomy in an ACL reconstructed population. However, in the study of Keays *et al*,³⁷ the relationship was close to significant and in another study meniscal injury/meniscectomy was significantly associated with patellofemoral OA.⁴¹ Furthermore, in a population without ACL injury, meniscectomy was related to development of patellofemoral OA.⁸⁵ An explanation for this relationship could be the influence of altered biomechanics in the knee, or the meniscal tear was a feature of the already existing early knee OA.

The results of this systematic review confirm the thoughts about the importance of preservation of the meniscus for preventing development of OA. Our advice for future studies is to document the location and extent of meniscectomy as well as which knee compartments, medial, lateral or patellofemoral, were used for assessing OA development.

Three key clinical questions and our findings

In clinical practice, three questions are important with regard to choice of treatment for ACL injuries and the development of knee OA.

1. *What is the influence of operative versus non-operative treatment on OA development?* On the basis of our results, we

cannot answer this question because there was conflicting evidence. However, we should note that, in the operatively treated patients, the graft type was mostly bone-patellar tendon-bone.^{30 31 40 41 47 53 65 70 72 77 79 80} So, there is less information on hamstring tendon reconstructed patients versus non-operatively treated patients and development of OA, despite both graft types being commonly used for ACL reconstruction.⁸⁶

2. *When operative treatment is chosen, what is the influence of graft choice?* On the basis of this systematic review, we cannot recommend one graft type to reduce OA risk.
3. *Is early reconstruction necessary for preventing OA development?* The aim of early timing of reconstruction after ACL rupture is to prevent new meniscal and cartilage damage. Our results indicate that early or late reconstruction is not related to greater risk of patellofemoral or tibiofemoral OA.

However, for OA development in unspecified compartment, we cannot give any indication which time point, early or late after injury, is best for reconstruction with regard to preventing OA development. A possible explanation for these conflicting results is the heterogeneity of additional injuries in the included studies and differences in the definition of early reconstruction. Furthermore, Smith *et al*⁸⁷ found in their meta-analysis no significant difference in the incidence of chondral and meniscal injuries between early and delayed reconstruction groups (the latter was defined as a minimum of 6 weeks postinjury). Another explanation might be that degenerative changes develop after the initial trauma caused by, for example, traumatic bone marrow lesions and activation of proinflammatory cytokines, independently of the choice of treatment.³ Besides, ACL reconstruction is a new trauma with additional damage such as bone marrow lesions, haemarthrosis and inflammation-related factors, for example, inflammatory cytokines.

Other considerations

We did not distinguish between partial and complete ACL tears. Partial or complete tears need to be diagnosed by arthroscopic evaluation, the reference for diagnosing ACL rupture. We may assume that the studies that included operatively treated patients enrolled patients with complete ACL tears. However, most studies did not describe their arthroscopic findings. Of the four studies which included non-operatively treated patients, one⁶⁴ reported the inclusion of both partial and complete tears, two^{22 76} reported the inclusion of only complete tears and one⁷¹ did not describe the type of the ACL tear. Thus, it is difficult to draw conclusions about the difference between the influence of partial and complete tears on OA development. Besides, in long-term follow-up studies, it is possible that partial tears progress to complete ACL tears⁸⁸ and then it is difficult to distinguish the contribution of the partial and complete tears to the development of OA.

A determinant, which was not included in the results, is the altered knee biomechanics after ACL injury. The possible explanation for no information about this determinant is that studies researching the altered knee biomechanics include fewer patients ($n \leq 20$, exclusion criteria of this systematic review) and that these studies have a cross-sectional design (exclusion criteria of this systematic review). Chaudari *et al*⁸⁹ suggest that the observed changes in the knee biomechanics result in altered loading patterns and influence metabolic changes in the underlying cartilage. Reduced internal tibial rotation was found in patients after ACL reconstruction compared with the contralateral knee and healthy controls.⁹⁰ In addition to this finding, a recently published cross-sectional study showed that after ACL reconstruction, patients with patellofemoral OA and valgus

alignment had significantly less internal knee rotation during walking and running than patients with valgus alignment and no patellofemoral OA.⁹¹ However, this study had a cross-sectional design; prospective studies are required to evaluate if the altered knee rotation is a result of patellofemoral OA or influences the development of patellofemoral OA.

Limitations

This systematic review has some limitations. First, of the 64 included studies, only 14^{23 26 34 35 38 42 43 46 47 56 65 67 68 70} corrected for the influence of confounders. Consequently, the reported influence of determinants on the development of OA may be partly or completely explained by other factors. By presenting the data, one of the criteria to be classified as a low risk of bias study was controlling for confounding. A prospective observational study design is the best way to determine predictors for development of OA after an ACL rupture. However, prospective collected data and retrospective analyses (research question defined after data collection) were also useful for our research question. Therefore, we also included retrospective study designs.

Second, the number of patients available for analysis at follow-up in the included studies was small. Only 18 of the 64 (28%) included studies had more than 100 patients available for analysis at follow-up.

Third, the included studies were heterogeneous with regard to study design, determinant assessment, additional intra-articular injuries, reported OA outcome, definition of OA and the statistical methods used. For these reasons, comparison between the included studies was difficult and pooling of the data was not possible. Therefore, we used the second best option for presenting the results: best-evidence synthesis.

Best-evidence synthesis is appropriate for summarising the available evidence. All the 64 included studies were classified as low risk or high risk of bias; however, only two studies met the criteria for low risk of bias. This means that reporting of inclusion of consecutive patients, measurement of determinants and outcomes independently, using accurate measures for the determinants and description of loss to follow-up with a maximal 20% and correction for confounding were poorly performed and described in the included studies.

Finally, we attempted to evaluate the influence of determinants on the development of tibiofemoral and patellofemoral OA separately. However, we should note that some studies did not use a valid tool for the compartmental assessment of OA, (eg, Kellgren and Lawrence score for assessment of patellofemoral OA). In some studies, the compartment was not described (compartment unspecified). The evaluation of the correctly used classification system for compartmental OA assessment was not included in the quality assessment tool.

Strengths

The strengths of this systematic review are that we summarised the evidence for tibiofemoral OA and patellofemoral OA outcomes after ACL injury separately. Moreover, we summarised these outcomes in patients who had had ACL reconstruction and those who had been managed with conservative treatment. Additionally, we evaluated determinants that influence early degenerative changes because we included studies with relatively short follow-ups (a minimum of 2 years). To be comprehensive, we chose to include both prospective and retrospective study designs having at least 20 patients. In addition to previously published systematic reviews,⁵⁻⁷ we included 21 studies published after the search dates of those systematic reviews.

Studies that used outdated surgery techniques were excluded, which resulted in exclusion of many older studies. However, our oldest study included was published in 1989⁶⁴ and newer studies might be of better quality as our two low risk of bias studies were published in 2012⁵⁶ and 2013.³⁵ The best-evidence synthesis considers the quality of the studies and accounts for a possible bias. When we analysed the results of studies only published during the past 10 years, the results differed minimally. The only aspects that changed were the influence of chondral injury (location not reported) on OA development (compartment unspecified), and of the graft type bone-patellar tendon-bone; both would change from conflicting evidence to limited evidence for a positive relationship with development of OA. These results of limited evidence still need more high-quality studies in order to make firm recommendations.

Overall, we can conclude that despite the inclusion of many new studies in this comprehensive systematic review, including two low risk of bias studies,^{35 56} more low risk of bias studies are required to evaluate determinants and their role in OA development. Many determinants showed conflicting and limited evidence. The following determinants should be further studied in large prospective studies, which could be used for meta-analysis: knee function and activity level, both examined in the first period after ACL rupture, patients characteristics, such as age, BMI and sex, meniscal injury/menisectomy specified in medial and lateral compartments, meniscus repair, chondral injury, choice of treatment, graft type and reconstruction technique. We strongly recommend specifying the compartment of OA development.

In summary, medial meniscal injury/menisectomy after ACL rupture influences the development of OA (tibiofemoral OA and compartment unspecified). In contrast, it seems that lateral meniscal injury/menisectomy has no relationship with OA development. Our results also suggest that time between injury and reconstruction does not influence the development of patellofemoral and tibiofemoral OA. However, we found limited or conflicting evidence for many determinants.

What are the new findings?

In patients with an anterior cruciate ligament rupture:

- Moderate evidence was found that medial meniscus injury/menisectomy had influence on osteoarthritis (OA) development; in contrast, lateral meniscus injury/menisectomy showed moderate evidence for no relationship with development of OA.
- Time between injury and reconstruction showed moderate evidence for no relationship with patellofemoral and tibiofemoral OA development.
- It is still unclear which treatment option is the best for preventing OA development; conflicting evidence was found between treatment choice (operative vs non-operative treatment) and development of knee OA.

Acknowledgements The authors would like to thank Louis Volkers for his help with the literature search.

Contributors All authors were involved in designing this systematic review, interpreting the data and have contributed to the writing and editing of the manuscript. BLvM and MR assessed the studies for inclusion. BLvM and WAvE performed the data extraction. MR, SMAB-Z, DEM, JANV and BLvM performed the risk of bias assessment.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Moses B, Orchard J, Orchard J. Systematic review: annual incidence of ACL injury and surgery in various populations. *Res Sports Med* 2012;20:157–79.
- 2 Gillquist J, Messner K. Anterior cruciate ligament reconstruction and the long-term incidence of gonarthrosis. *Sports Med* 1999;27:143–56.
- 3 Lohmander LS, Englund PM, Dahl LL, et al. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med* 2007;35:1756–69.
- 4 Renström PA. Eight clinical conundrums relating to anterior cruciate ligament (ACL) injury in sport: recent evidence and a personal reflection. *Br J Sports Med* 2013;47:367–72.
- 5 Oiestad BE, Engebretsen L, Storheim K, et al. Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. *Am J Sports Med* 2009;37:1434–43.
- 6 Magnussen RA, Spindler KP. The effect of patient and injury factors on long-term outcome after anterior cruciate ligament reconstruction. *Curr Orthop Pract* 2011;22:90–103.
- 7 Claes S, Hermie L, Verdonk R, et al. Is osteoarthritis an inevitable consequence of anterior cruciate ligament reconstruction? A meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 2013;21:1967–76.
- 8 Culvenor AG, Cook JL, Collins NJ, et al. Is patellofemoral joint osteoarthritis an under-recognised outcome of anterior cruciate ligament reconstruction? A narrative literature review. *Br J Sports Med* 2013;47:66–70.
- 9 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
- 10 Altman R, Asch E, Bloch D, et al. Development of criteria for the classification and reporting of osteoarthritis. Classification of osteoarthritis of the knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism Association. *Arthritis Rheum* 1986;29:1039–49.
- 11 Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52:377–84.
- 12 Slim K, Nini E, Forestier D, et al. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003;73:712–16.
- 13 Deeks JJ, Dinnes J, D'Amico R, et al. Evaluating non-randomised intervention studies. *Health Technol Assess* 2003;7:iii–x, 1–173.
- 14 Guyatt GH, Sackett DL, Sinclair JC, et al. Users' guides to the medical literature. IX. A method for grading health care recommendations. Evidence-Based Medicine Working Group. *JAMA* 1995;274:1800–4.
- 15 Slavin RE. Best evidence synthesis: an intelligent alternative to meta-analysis. *J Clin Epidemiol* 1995;48:9–18.
- 16 van Tulder M, Furlan A, Bombardier C, et al. Updated method guidelines for systematic reviews in the Cochrane collaboration back review group. *Spine (Phila Pa 1976)* 2003;28:1290–9.
- 17 Furlan AD, van Tulder M, Cherkin D, et al. Acupuncture and dry-needling for low back pain: an updated systematic review within the framework of the Cochrane collaboration. *Spine (Phila Pa 1976)* 2005;30:944–63.
- 18 Ahliden M, Kartus J, Ejerhed L, et al. Knee laxity measurements after anterior cruciate ligament reconstruction, using either bone-patellar-tendon-bone or hamstring tendon autografts, with special emphasis on comparison over time. *Knee Surg Sports Traumatol Arthrosc* 2009;17:1117–24.
- 19 Frobell RB, Roos HP, Roos EM, et al. Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. *BMJ* 2013;346:f232.
- 20 Harilainen A, Linko E, Sandelin J. Randomized prospective study of ACL reconstruction with interference screw fixation in patellar tendon autografts versus femoral metal plate suspension and tibial post fixation in hamstring tendon autografts: 5-year clinical and radiological follow-up results. *Knee Surg Sports Traumatol Arthrosc* 2006;14:517–28.
- 21 Holm I, Oiestad BE, Risberg MA, et al. No difference in knee function or prevalence of osteoarthritis after reconstruction of the anterior cruciate ligament with 4-strand hamstring autograft versus patellar tendon-bone autograft: a randomized study with 10-year follow-up. *Am J Sports Med* 2010;38:448–54.
- 22 Meunier A, Odensten M, Good L. Long-term results after primary repair or non-surgical treatment of anterior cruciate ligament rupture: a randomized study with a 15-year follow-up. *Scand J Med Sci Sports* 2007;17:230–7.
- 23 O'Neill DB. Arthroscopically assisted reconstruction of the anterior cruciate ligament. A follow-up report. *J Bone Joint Surg Am* 2001;83-A:1329–32.
- 24 Sajovic M, Strahovnik A, Dernovsek MZ, et al. Quality of life and clinical outcome comparison of semitendinosus and gracilis tendon versus patellar tendon autografts for anterior cruciate ligament reconstruction: an 11-year follow-up of a randomized controlled trial. *Am J Sports Med* 2011;39:2161–9.
- 25 Song EK, Seon JK, Yim JH, et al. Progression of osteoarthritis after double- and single-bundle anterior cruciate ligament reconstruction. *Am J Sports Med* 2013;41:2340–6.

- 26 Sun K, Tian SQ, Zhang JH, *et al.* Anterior cruciate ligament reconstruction with bone-patellar tendon-bone autograft versus allograft. *Arthroscopy* 2009;25:750–9.
- 27 Suomalainen P, Jarvela T, Paakkala A, *et al.* Double-bundle versus single-bundle anterior cruciate ligament reconstruction: a prospective randomized study with 5-year results. *Am J Sports Med* 2012;40:1511–18.
- 28 Wipfler B, Donner S, Zechmann CM, *et al.* Anterior cruciate ligament reconstruction using patellar tendon versus hamstring tendon: a prospective comparative study with 9-year follow-up. *Arthroscopy* 2011;27:653–65.
- 29 Zaffagnini S, Bruni D, Marcheggiani Muccioli GM, *et al.* Single-bundle patellar tendon versus non-anatomical double-bundle hamstrings ACL reconstruction: a prospective randomized study at 8-year minimum follow-up. *Knee Surg Sports Traumatol Arthrosc* 2011;19:390–7.
- 30 Fink C, Hoser C, Hackl W, *et al.* Long-term outcome of operative or nonoperative treatment of anterior cruciate ligament rupture—is sports activity a determining variable? *Int J Sports Med* 2001;22:304–9.
- 31 Fithian DC, Paxton EW, Stone ML, *et al.* Prospective trial of a treatment algorithm for the management of the anterior cruciate ligament-injured knee. *Am J Sports Med* 2005;33:335–46.
- 32 Giron F, Aglietti P, Cuomo P, *et al.* Anterior cruciate ligament reconstruction with double-looped semitendinosus and gracilis tendon graft directly fixed to cortical bone: 5-year results. *Knee Surg Sports Traumatol Arthrosc* 2005;13:81–91.
- 33 Hanytsiak BT, Spindler KP, Rothrock CR, *et al.* Twelve-year follow-up on anterior cruciate ligament reconstruction: long-term outcomes of prospectively studied osseous and articular injuries. *Am J Sports Med* 2008;36:671–7.
- 34 Hui C, Salmon LJ, Kok A, *et al.* Fifteen-year outcome of endoscopic anterior cruciate ligament reconstruction with patellar tendon autograft for “isolated” anterior cruciate ligament tear. *Am J Sports Med* 2011;39:89–98.
- 35 Janssen RP, du Mee AW, van Valkenburg J, *et al.* Anterior cruciate ligament reconstruction with 4-strand hamstring autograft and accelerated rehabilitation: a 10-year prospective study on clinical results, knee osteoarthritis and its predictors. *Knee Surg Sports Traumatol Arthrosc* 2013;21:1977–88.
- 36 Jomha NM, Borton DC, Clingeleffer AJ, *et al.* Long-term osteoarthritic changes in anterior cruciate ligament reconstructed knees. *Clin Orthop Relat Res* 1999;188–93.
- 37 Keays SL, Newcombe PA, Bullock-Saxton JE, *et al.* Factors involved in the development of osteoarthritis after anterior cruciate ligament surgery. *Am J Sports Med* 2010;38:455–63.
- 38 Leys T, Salmon L, Waller A, *et al.* Clinical results and risk factors for reinjury 15 years after anterior cruciate ligament reconstruction: a prospective study of hamstring and patellar tendon grafts. *Am J Sports Med* 2012;40:595–605.
- 39 Moisala AS, Jarvela T, Honkonen S, *et al.* Arthroscopic anterior cruciate ligament reconstruction using a hamstring graft with interference screw fixation. *Scand J Surg* 2007;96:83–7.
- 40 Neuman P, Englund M, Kostogiannis I, *et al.* Prevalence of tibiofemoral osteoarthritis 15 years after nonoperative treatment of anterior cruciate ligament injury: a prospective cohort study. *Am J Sports Med* 2008;36:1717–25.
- 41 Neuman P, Kostogiannis I, Friden T, *et al.* Patellofemoral osteoarthritis 15 years after anterior cruciate ligament injury—a prospective cohort study. *Osteoarthritis Cartilage* 2009;17:284–90.
- 42 Oiestad BE, Holm I, Engebretsen L, *et al.* The prevalence of patellofemoral osteoarthritis 12 years after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2013;21:942–9.
- 43 Oiestad BE, Holm I, Gunderson R, *et al.* Quadriceps muscle weakness after anterior cruciate ligament reconstruction: a risk factor for knee osteoarthritis? *Arthritis Care Res (Hoboken)* 2010;62:1706–14.
- 44 Oiestad BE, Holm I, Aune AK, *et al.* Knee function and prevalence of knee osteoarthritis after anterior cruciate ligament reconstruction: a prospective study with 10 to 15 years of follow-up. *Am J Sports Med* 2010;38:2201–10.
- 45 Pinczewski LA, Lyman J, Salmon LJ, *et al.* A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and patellar tendon autograft: a controlled, prospective trial. *Am J Sports Med* 2007;35:564–74.
- 46 Pinczewski LA, Salmon LJ, Jackson WF, *et al.* Radiological landmarks for placement of the tunnels in single-bundle reconstruction of the anterior cruciate ligament. *J Bone Joint Surg Br* 2008;90:172–9.
- 47 Potter HG, Jain SK, Ma Y, *et al.* Cartilage injury after acute, isolated anterior cruciate ligament tear: immediate and longitudinal effect with clinical/MRI follow-up. *Am J Sports Med* 2012;40:276–85.
- 48 Ruiz AL, Kelly M, Nutton RW. Arthroscopic ACL reconstruction: a 5–9 year follow-up. *Knee* 2002;9:197–200.
- 49 Shelbourne KD, Gray T. Results of anterior cruciate ligament reconstruction based on meniscus and articular cartilage status at the time of surgery. Five- to fifteen-year evaluations. *Am J Sports Med* 2000;28:446–52.
- 50 Shelbourne KD, Urch SE, Gray T, *et al.* Loss of normal knee motion after anterior cruciate ligament reconstruction is associated with radiographic arthritic changes after surgery. *Am J Sports Med* 2012;40:108–13.
- 51 Wu WH, Hackett T, Richmond JC. Effects of meniscal and articular surface status on knee stability, function, and symptoms after anterior cruciate ligament reconstruction: a long-term prospective study. *Am J Sports Med* 2002;30:845–50.
- 52 Mascarenhas R, Tranovich MJ, Kropf EJ, *et al.* Bone-patellar tendon-bone autograft versus hamstring autograft anterior cruciate ligament reconstruction in the young athlete: a retrospective matched analysis with 2–10 year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2012;20:1520–7.
- 53 Meuffels DE, Favejee MM, Visser MM, *et al.* Ten year follow-up study comparing conservative versus operative treatment of anterior cruciate ligament ruptures. A matched-pair analysis of high level athletes. *Br J Sports Med* 2009;43:347–51.
- 54 Aglietti P, Zaccherotti G, De Biase P, *et al.* A comparison between medial meniscus repair, partial meniscectomy, and normal meniscus in anterior cruciate ligament reconstructed knees. *Clin Orthop Relat Res* 1994:165–73.
- 55 Aglietti P, Buzzi R, Giron F, *et al.* Arthroscopic-assisted anterior cruciate ligament reconstruction with the central third patellar tendon. A 5–8-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 1997;5:138–44.
- 56 Ahn JH, Kim JG, Wang JH, *et al.* Long-term results of anterior cruciate ligament reconstruction using bone-patellar tendon-bone: an analysis of the factors affecting the development of osteoarthritis. *Arthroscopy* 2012;28:1114–23.
- 57 Cohen M, Amaro JT, Eijnsman B, *et al.* Anterior cruciate ligament reconstruction after 10 to 15 years: association between meniscectomy and osteoarthritis. *Arthroscopy* 2007;23:629–34.
- 58 Gerhard P, Bolt R, Duck K, *et al.* Long-term results of arthroscopically assisted anatomical single-bundle anterior cruciate ligament reconstruction using patellar tendon autograft: are there any predictors for the development of osteoarthritis? *Knee Surg Sports Traumatol Arthrosc* 2013;21:957–64.
- 59 Hart AJ, Buscombe J, Malone A, *et al.* Assessment of osteoarthritis after reconstruction of the anterior cruciate ligament: a study using single-photon emission computed tomography at ten years. *J Bone Joint Surg Br* 2005;87:1483–7.
- 60 Ichiba A, Kishimoto I. Effects of articular cartilage and meniscus injuries at the time of surgery on osteoarthritic changes after anterior cruciate ligament reconstruction in patients under 40 years old. *Arch Orthop Trauma Surg* 2009;129:409–15.
- 61 Jarvela T, Nyyssönen M, Kannus P, *et al.* Bone-patellar tendon-bone reconstruction of the anterior cruciate ligament. A long-term comparison of early and late repair. *Int Orthop* 1999;23:227–31.
- 62 Jarvela T, Paakkala T, Kannus P, *et al.* The incidence of patellofemoral osteoarthritis and associated findings 7 years after anterior cruciate ligament reconstruction with a bone-patellar tendon-bone autograft. *Am J Sports Med* 2001;29:18–24.
- 63 Jarvela T, Kannus P, Jarvinen M. Anterior cruciate ligament reconstruction in patients with or without accompanying injuries: a re-examination of subjects 5 to 9 years after reconstruction. *Arthroscopy* 2001;17:818–25.
- 64 Kannus P, Jarvinen M. Posttraumatic anterior cruciate ligament insufficiency as a cause of osteoarthritis in a knee joint. *Clin Rheumatol* 1989;8:251–60.
- 65 Kessler MA, Behrend H, Henz S, *et al.* Function, osteoarthritis and activity after ACL-rupture: 11 years follow-up results of conservative versus reconstructive treatment. *Knee Surg Sports Traumatol Arthrosc* 2008;16:442–8.
- 66 Lebel B, Hulet C, Galaud B, *et al.* Arthroscopic reconstruction of the anterior cruciate ligament using bone-patellar tendon-bone autograft: a minimum 10-year follow-up. *Am J Sports Med* 2008;36:1275–82.
- 67 Leiter JR, Goulay R, McRae S, *et al.* Long-term follow-up of ACL reconstruction with hamstring autograft. *Knee Surg Sports Traumatol Arthrosc* 2014;22:1061–9.
- 68 Li RT, Lorenz S, Xu Y, *et al.* Predictors of radiographic knee osteoarthritis after anterior cruciate ligament reconstruction. *Am J Sports Med* 2011;39:2595–603.
- 69 Liden M, Sernert N, Rostgard-Christensen L, *et al.* Osteoarthritic changes after anterior cruciate ligament reconstruction using bone-patellar tendon-bone or hamstring tendon autografts: a retrospective, 7-year radiographic and clinical follow-up study. *Arthroscopy* 2008;24:899–908.
- 70 Lohmander LS, Ostergren A, Englund M, *et al.* High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum* 2004;50:3145–52.
- 71 Menke W, Schneider T, Schmitz B, *et al.* [Results of follow-up of untreated anterior cruciate ligament rupture] Nachuntersuchungsergebnisse bei unbehandelter vorderer Kreuzbandruptur. *Sportverletz Sportschaden* 1990;4:169–74.
- 72 Mihelic R, Jurdana H, Jotanovic Z, *et al.* Long-term results of anterior cruciate ligament reconstruction: a comparison with non-operative treatment with a follow-up of 17–20 years. *Int Orthop* 2011;35:1093–7.
- 73 Murray JR, Lindh AM, Hogan NA, *et al.* Does anterior cruciate ligament reconstruction lead to degenerative disease? Thirteen-year results after bone-patellar tendon-bone autograft. *Am J Sports Med* 2012;40:404–13.
- 74 Otto D, Pinczewski LA, Clingeleffer A, *et al.* Five-year results of single-incision arthroscopic anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 1998;26:181–8.
- 75 Salmon LJ, Russell VJ, Refshauge K, *et al.* Long-term outcome of endoscopic anterior cruciate ligament reconstruction with patellar tendon autograft: minimum 13-year review. *Am J Sports Med* 2006;34:721–32.
- 76 Segawa H, Omori G, Koga Y. Long-term results of non-operative treatment of anterior cruciate ligament injury. *Knee* 2001;8:5–11.
- 77 Seitz H, Chrysopoulos A, Egkher E, *et al.* [Long-term results of replacement of the anterior cruciate ligament in comparison with conservative therapy]

- Langzeitergebnisse nach vorderem Kreuzbandersatz im Vergleich zur konservativen Therapie. *Chirurg* 1994;65:992–8.
- 78 Seon JK, Song EK, Park SJ. Osteoarthritis after anterior cruciate ligament reconstruction using a patellar tendon autograft. *Int Orthop* 2006;30:94–8.
- 79 Streich NA, Zimmermann D, Bode G, *et al.* Reconstructive versus non-reconstructive treatment of anterior cruciate ligament insufficiency. A retrospective matched-pair long-term follow-up. *Int Orthop* 2011;35:607–13.
- 80 von Porat A, Roos EM, Roos H. High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. *Ann Rheum Dis* 2004;63:269–73.
- 81 Wang CJ, Huang TW, Jih S. Radiographic assessment of the knee after patellar tendon reconstruction for chronic anterior cruciate ligament deficiency. *Chang Gung Med J* 2004;27:85–90.
- 82 Thompson WO, Thaete FL, Fu FH, *et al.* Tibial meniscal dynamics using three-dimensional reconstruction of magnetic resonance images. *Am J Sports Med* 1991;19:210–15; discussion 5–6.
- 83 Salata MJ, Gibbs AE, Sekiya JK. A systematic review of clinical outcomes in patients undergoing meniscectomy. *Am J Sports Med* 2010;38:1907–16.
- 84 Crossley KM. Is patellofemoral osteoarthritis a common sequela of patellofemoral pain? *Br J Sports Med* 2014;48:409–10.
- 85 Englund M, Lohmander LS. Patellofemoral osteoarthritis coexistent with tibiofemoral osteoarthritis in a meniscectomy population. *Ann Rheum Dis* 2005;64:1721–6.
- 86 Li S, Chen Y, Lin Z, *et al.* A systematic review of randomized controlled clinical trials comparing hamstring autografts versus bone-patellar tendon-bone autografts for the reconstruction of the anterior cruciate ligament. *Arch Orthop Trauma Surg* 2012;132:1287–97.
- 87 Smith TO, Davies L, Hing CB. Early versus delayed surgery for anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 2010;18:304–11.
- 88 Noyes FR, Moar LA, Moorman CT III, *et al.* Partial tears of the anterior cruciate ligament. Progression to complete ligament deficiency. *J Bone Joint Surg Br* 1989;71:825–33.
- 89 Chaudari AM, Briant PL, Bevil SL, *et al.* Knee kinematics, cartilage morphology, and osteoarthritis after ACL injury. *Med Sci Sports Exerc* 2008;40:215–22.
- 90 Webster KE, Feller JA. Alterations in joint kinematics during walking following hamstring and patellar tendon anterior cruciate ligament reconstruction surgery. *Clin Biomech* 2011;26:175–80.
- 91 Culvenor AG, Schache AG, Vicenzino B, *et al.* Are knee biomechanics different in those with and without patellofemoral osteoarthritis after anterior cruciate ligament reconstruction? *Arthritis Car Res* 2014;66:1566–70.