

# One small step for man, one giant leap for men's health: a meta-analysis of behaviour change interventions to increase men's physical activity

Paul Sharp <sup>1</sup>, John C Spence,<sup>2</sup> Joan L Botorff,<sup>3,4</sup> John L Oliffe,<sup>5</sup> Kate Hunt,<sup>6</sup> Mathew Vis-Dunbar,<sup>7</sup> Cristina M Caperchione<sup>1,3</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2019-100912>).

<sup>1</sup>Faculty of Health, University of Technology Sydney, Sydney, New South Wales, Australia

<sup>2</sup>Sedentary Living Lab, Faculty of Kinesiology, Sport, and Recreation, University of Alberta, Edmonton, Alberta, Canada

<sup>3</sup>Institute for Healthy Living and Chronic Disease Prevention, University of British Columbia, Kelowna, British Columbia, Canada

<sup>4</sup>School of Nursing, University of British Columbia, Kelowna, British Columbia, Canada

<sup>5</sup>School of Nursing, University of British Columbia, Vancouver, British Columbia, Canada

<sup>6</sup>Institute for Social Marketing, Faculty of Health Sciences and Sport, University of Stirling, Stirling, UK

<sup>7</sup>Library, University of British Columbia, Kelowna, British Columbia, Canada

## Correspondence to

Paul Sharp, Human Performance Research Centre, Faculty of Health, University of Technology Sydney, Sydney, NSW 2007, Australia; [paul.sharp@uts.edu.au](mailto:paul.sharp@uts.edu.au)

Accepted 16 January 2020

Published Online First 5 February 2020

## ABSTRACT

**Objective** To determine the effects of behaviour change interventions on men's physical activity (postintervention), sustained change in physical activity behaviour ( $\geq 12$  months postintervention) and to identify variations in effects due to potential moderating variables (eg, theoretical underpinning, gender-tailored, contact frequency).

**Design** Systematic review with meta-analysis. Pooled effect size (Cohen's  $d$ ) was calculated assuming a random-effects model. Homogeneity and subsequent exploratory moderator analyses were assessed using  $Q$ ,  $T^2$  and  $I^2$ .

**Data sources** Medline, EMBASE, CINAHL, SportDiscus and Web of Science to April 2019.

**Eligibility criteria for selected studies** Randomised control trials of behaviour change interventions in men ( $\geq 18$  years) where physical activity was an outcome and data were from men-only studies or disaggregated by sex.

**Results** Twenty-six articles described 24 eligible studies. The overall mean intervention effect on men's physical activity was 0.35 (SE=0.05; 95% CI 0.26 to 0.45;  $p<0.001$ ). This effect size is consistent with an increase of approximately 97 min of total physical activity per week or 980 steps per day. Intervention moderators associated with greater increases in physical activity included objective physical activity outcome measures, a gender-tailored design, use of a theoretical framework, shorter length programmes ( $\leq 12$  weeks), using four or more types of behaviour change techniques and frequent contact with participants ( $\geq 1$  contact per week). 12 studies included additional follow-up assessments ( $\geq 12$  months postintervention) and the overall mean effect was 0.32 (SE=0.09; 95% CI 0.15 to 0.48;  $p<0.001$ ) for that sustained increase in physical activity.

**Summary** Behaviour change interventions targeting men's physical activity can be effective. Moderator analyses are preliminary and suggest research directions.

## INTRODUCTION

Physical activity is important in disease prevention and illness management and there has been considerable research into effective physical activity interventions.<sup>1,2</sup> Although gender is recognised as an important sociocultural factor influencing health and health-related behaviours,<sup>3</sup> its influence on the uptake of behavioural interventions is not well understood.

Worldwide, women live for almost 6 years longer than men and men have higher rates of all-cause

mortality for many conditions.<sup>4</sup> Factors associated with these sex differences are men's alignments to health compromising, masculine roles, identities and relations.<sup>5</sup> These expressions of masculinities intersect with other social determinants of health (eg, socioeconomic status, race) to marginalise some subgroups of men, creating significant health inequities. Consequently, some men lack the knowledge and/or resources to promote their health and/or access health services and may be less willing to attend health education sessions than women.<sup>6–9</sup> It is often assumed that men cannot or will not access health promotion programmes, and that programmes designed for the general public will suffice for those men who are willing to attend.

Evaluations of health promotion programmes often fall short in providing information about effective strategies to promote men's health because of the under representation of men. For example, in a meta-analysis demonstrating a small effect for adult physical activity interventions ( $d=0.19$ ), only 26% of participants within the included studies were men,<sup>2</sup> making generalisability challenging. As approximately 31% of the population worldwide is insufficiently active,<sup>10</sup> and clinically relevant health benefits may be accrued through relatively small increases in physical activity,<sup>11</sup> millions of men stand to benefit from effective health promotion interventions.

To yield the full health benefits of being physically active, behaviour must be sustained over time. A limited number of studies include additional follow-up measures postintervention and the evidence for long-term behaviour change is mixed.<sup>12–14</sup> A Cochrane review investigating the effectiveness of interventions for promoting self-reported physical activity found that intervention effects postintervention ( $d=0.28$ ) were not maintained in 6 of the 19 studies reporting outcomes after 6 months.<sup>12</sup> Two more recent reviews examining the effects of web-based and face-to-face interventions reported small effects at 12 months postintervention ( $d=0.20$ , 0.19, respectively) and either no or small effects at 24 months. However, most studies failed to measure the long-term (ie,  $\geq 12$  months) effects on physical activity.<sup>13,14</sup>

Developments in men's health promotion have resulted in an increased number of physical activity interventions targeted at engaging and retaining men.<sup>15,16</sup> Several strategies and approaches (eg, men-only programmes, 'masculine' setting) have been identified that show promise for improving men's



© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Sharp P, Spence JC, Botorff JL, et al. *Br J Sports Med* 2020;**54**:1208–1216.

participation, retention and overall success rates.<sup>15–20</sup> Interventions tailored specifically to the values, preferences and interests of men (eg, gender-tailored) may increase programme effectiveness.<sup>15</sup> Despite the growing interest in men's health promotion, the overall effectiveness of physical activity interventions for men remains unclear. The aim of our meta-analysis was to determine the effects of behaviour change interventions to increase men's physical activity, sustained physical activity change ( $\geq 12$  months postintervention) and to identify how potential moderating variables were associated with key outcomes (eg, theoretical underpinning, gender-tailored, contact frequency).

## METHODS

This review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) Statement (online supplementary table S1)<sup>21</sup> and was prospectively registered in the PROSPERO registry of systematic reviews (#CRD42018079448).

### Inclusion criteria

Eligible studies for inclusion were randomised controlled trials (RCTs) identified using the following framework:

1. **Population.** Studies included adult men age 18+ years. Mixed sex studies were included provided relevant data (see outcomes) for men were reported separately. Consistent with previous research,<sup>15 16</sup> studies that exclusively included older adults ( $\geq 65$  years) were excluded as they are likely to have different intervention requirements.
2. **Intervention.** Interventions with clear and deliberate intent to increase the physical activity levels of participants. Physical activity was defined as any bodily movement that increased energy expenditure beyond basal levels.<sup>2</sup> Diverse physical activity behaviour change interventions were eligible (eg, education sessions, supervised physical activity practice sessions). Articles that included both physical activity and other health behaviours (eg, diet) were included, provided that physical activity change was an intended and explicitly reported outcome.
3. **Comparison.** Studies were RCTs.
4. **Outcomes.** An outcome measure of physical activity (eg, steps per day, total activity minutes per week), disaggregated by sex (if applicable), available for both intervention and control groups, representing physical activity change from prepoint to postpoint or multipoint test.

### Search method

A comprehensive search strategy was undertaken to identify all possible studies for inclusion. The search was applied to MEDLINE and adapted for EMBASE, CINAHL, SportDiscus and Web of Science. All searches were completed by a specialised research librarian (MVD) to April 2019. Search terms included MeSH and keywords relevant to the aims and in accordance with the eligibility criteria: (1) population (eg, Male/ or (men or male?)); (2) intervention (eg, Exercise/ or ("physical activit\*" or exercis\*)) and (3) outcomes (eg, Fitness Trackers/ or Self Report/). Additional filters were used to limit results to RCTs (eg, Randomized Controlled Trial/ or (randomi#ed or experimental) adj3 trial). Searches were limited to English language, original research and academic journals. No editorials, reviews, commentaries, conference abstracts or other grey literature were included. The decision to not include grey literature was based on concerns relating to the absence of peer-review and the potential for identifying an unrepresentative sample of all

unpublished studies. The reference lists of included articles were manually searched for potential studies not yet identified. See online supplementary table S2 for the complete search syntax. Prior to manuscript submission, identified articles were reviewed to ensure that no trial had been retracted between inclusion and publication.<sup>22</sup>

### Screening of articles

All identified references were imported into EndNote X8 (Clarivate Analytics, Philadelphia, Pennsylvania, USA). Duplicates were automatically identified by matches in authorship, year and title and manually reviewed prior to deletion. Overseen by the lead author (PS), two trained research assistants performed a title and abstract review to screen remaining records for relevance. Full text articles were retrieved for all remaining records and further screened to identify the final set of articles for inclusion. Any uncertainty was discussed among the research team.

### Data extraction, study quality and quality of evidence

A coding framework was developed, pilot tested and refined by two researchers (PS and JCS). Study characteristics were coded by two reviewers (PS and JCS) under four general categories relating to the study design (eg, sample size, physical activity measurement), participants (eg, mean age, health status), intervention (eg, mode of delivery, behaviour change techniques, theoretical underpinning) and results (eg, mean change, SD). Outcome data for use in the meta-analysis were recorded for baseline, immediately postintervention and 12 months or greater postintervention. If more than one variable was available for physical activity, the variable that best reflected an overall measure of physical activity was selected (eg, total MET-minutes, self-reported total physical activity). Interventions were coded using Michie and colleagues<sup>23</sup> definitions for characterising behaviour change interventions including education, persuasion, incentivisation, coercion, training, restriction, environmental restructuring, modelling and enablement. Relevant detail was sought from additional publications (eg, protocol papers), when available. Interventions were deemed to be gender-tailored if there was evidence to suggest that they were designed specifically to the values, preferences and interests of men. Intervention engagement was assessed as high ( $>80\%$ ), moderate ( $60\%–80\%$ ) or low ( $<60\%$ ) based on participants' average reported uptake of the intervention content (eg, attendance, website visits) or the extent to which participants met reported engagement goals (eg, % attending 10 of 12 sessions). The coding framework is not exhaustive of all intervention aspects and only common comparable characteristics reported in sufficient detail across studies are subsequently reported on.

Study quality was independently assessed by two members of the research team (PS and CMC) using The Effective Public Health Practice Project (EPHPP) tool.<sup>24</sup> This tool has been reported to have content and construct validity, excellent inter-rater reliability and is recommended by the Cochrane Public Health Review Group for assessing the quality of public health and health promotion studies.<sup>25–27</sup> This six-domain (14 question) rating scale for interventions assesses selection bias, study design, assessment of confounders, data collection methods (reliability and validity) and reporting of blinding, withdrawals and dropouts. In accordance with the tool's guidelines, a score for each domain of weak (1 point), moderate (2 points) or strong (3 points) was awarded and averaged to provide a total score for each study. Based on their total score, studies are assigned a quality rating of weak (1.00–1.50), moderate (1.51–2.50)


or strong (2.51–3.00). Where discrepancies existed between reviewers, deliberation occurred until consensus was reached.

Additionally, the overall quality of evidence was assessed by three members of the research team (PS, CMC and JLB) using GRADE.<sup>28 29</sup> The quality of evidence was performed for each study outcome (ie, physical activity change and long-term physical activity change) and reflects the extent to which we are confident that an estimate of the effect is correct. The quality of evidence can be assessed as high, moderate, low or very low. As all studies were randomised trials, study quality is initially assumed to be high but can be rated down based on risk of bias, inconsistency of results, indirectness of evidence, imprecision or publication bias.

### Statistical methods

Standardised mean differences (ie, effect size) with 95% CIs were computed to represent the effect of the interventions on men's physical activity. A positive effect size indicates a more favourable change in physical activity for the intervention condition. Cohen's criteria were used for interpretation of effect sizes as small (<0.50), moderate (0.50–0.79) and large (>0.79).<sup>30</sup> In addition, estimates of mean physical activity effect sizes were converted to the original metrics of ambulatory steps per day and minutes per week. Effect sizes were calculated using change from baseline scores, as this method removes a component of between-person variability from the analysis by controlling for preintervention differences.<sup>31</sup> In some cases, the required statistics were not reported. If available, and if possible, change scores were calculated from pretest and post-test means and SD, means and SEs, CI or other statistics (eg, p values), using conventional methods detailed by Borenstein *et al.*<sup>32</sup> In studies that included multiple intervention groups, a pooled mean and SD was calculated to create a single pair-wise comparison before entering the meta-analysis. Cluster-randomised trials were adjusted for using an estimation of the sample size.<sup>33</sup> In such instances, a design effect was calculated for each study using an intracluster correlation coefficient of 0.05, which has been previously used in meta-analyses of physical activity trials.<sup>34–37</sup>

Comprehensive meta-analysis V3 software was used for all analyses. A random effects model with inverse variance weighting was applied to estimate the pooled effect for physical activity. Two overall effect size calculations were conducted to investigate the effect of interventions on men's physical activity (baseline to postintervention) and long-term physical activity change (baseline to 12 month or greater postintervention). Each study contributed one effect size calculation to the overall analysis and 12 studies reported an additional follow-up measure 12 months or greater postintervention. Sensitivity analysis was conducted using the 'one-study remove' procedure. Publication bias was analysed using Egger's regression test,<sup>38</sup> Duval and Tweedie's Trim and Fill Procedure<sup>39</sup> and Rosenthal's Fail-Safe N calculation.<sup>40</sup> Homogeneity of effects was assessed through the Q-statistic. A significant Q-within ( $Q_w$ ) value indicates a heterogeneous distribution and suggests a need to conduct follow-up moderator analyses. To interpret heterogeneity, Tau-squared ( $T^2$ ), an estimate of total variance between studies, and I-squared ( $I^2$ ), a ratio of excess dispersion to total dispersion, were calculated. Larger  $T^2$  values reflect the proportion of variance that can be attributed to real differences between studies.  $I^2$  can be understood as the overlap of CIs explaining the total variance attributed to the covariates, interpreted as low (25%), moderate (50%) and high (75%) relative variance. Larger  $I^2$  values require



**Figure 1** Pathway of articles identified and excluded. PA, physical activity; RCT, randomised controlled trial.

techniques (ie, moderator analysis or meta-regression) to provide explanations.

Moderator analyses were conducted to explore potential variations in effectiveness due to differences in study, participant or interventions characteristics, using mixed effects analysis. Subgroup analyses were used to explore heterogeneity and make comparisons between characteristics. Categories were determined based on previous literature as well as the cut points that may be relevant for future intervention design.<sup>41</sup> A common among-study variance was assumed across subgroups and a pooled within-group estimate of  $T^2$  was used. In light of previous research exploring the effects of physical activity interventions<sup>42</sup> as well as work done in the field of men's health,<sup>15 16 19</sup> it can be reasonably argued that more intensive interventions (eg, greater contact frequency, using more types of behaviour change techniques) and interventions designed specifically for the target population (eg, gender-tailored) will be more effective. However, because previous research has provided a limited foundation for confirmatory hypothesis testing, the moderator analyses were considered exploratory and intended to be hypothesis-generating. Moderator analyses were not conducted for long-term physical activity change as only 12 studies included an additional follow-up 12 months or greater postintervention.

## RESULTS

### Description of included studies

The initial search strategy (excluding duplicates) identified 13 131 potentially relevant articles. Following title/abstract screening, 284 references remained from which an additional 258 articles were further removed following a full text review. Ultimately, 26 peer-reviewed journal articles (figure 1), representing 24 studies and independent samples,<sup>43–68</sup> were included in the review.

Tables 1 and 2 outline study design and intervention characteristics, respectively. Online supplementary table S3 provides an overview of the included interventions and additional details



Table 1 Study design and participant characteristics

Primary source	Study			Participant			
	Location	Design	PA measure	PA outcome*	N†	Mean age‡ (years)	Health status
Aguiar <i>et al</i> , 2016 <sup>43</sup>	Australia	RCT	Obj	Secondary	101	52	Overweight
Andersen <i>et al</i> , 2012 <sup>43 45</sup>	Norway	RCT	Obj	Primary	150	37	Inactive
Ashton <i>et al</i> , 2017 <sup>46</sup>	Australia	RCT	Obj	Primary	50	22	Inactive
Galvão <i>et al</i> , 2017 <sup>47</sup>	Australia	RCT	Sub	Primary	463	64	Cancer
Gong <i>et al</i> , 2015 <sup>48</sup>	China	Cluster	Sub	Secondary	450	64	Hypertension
Gray <i>et al</i> , 2013 <sup>49</sup>	Scotland	RCT	Sub	Secondary	103	47	Overweight
Groeneveld <i>et al</i> , 2011 <sup>50</sup>	Netherlands	RCT	Sub	Not clear	816	47	General
Hunt <i>et al</i> , 2014 <sup>51</sup>	Scotland	RCT	Sub	Secondary	747	47	Overweight
Livingston <i>et al</i> , 2015 <sup>52 53</sup>	Australia	Cluster	Sub	Primary	147	66	Cancer
Maruyama <i>et al</i> , 2010 <sup>54</sup>	Japan	RCT	Obj	Primary	110	40	MetS
McGowan <i>et al</i> , 2013 <sup>55</sup>	Canada	RCT	Sub	Primary	423	68	Cancer
Morgan <i>et al</i> , 2013 <sup>56</sup>	Australia	RCT	Obj	Secondary	159	48	Overweight
Morgan <i>et al</i> , 2014 <sup>57</sup>	Australia	RCT	Obj	Secondary	93	40	Overweight
Morgan <i>et al</i> , 2011a <sup>58</sup>	Australia	RCT	Obj	Secondary	53	41	Overweight
Morgan <i>et al</i> , 2011b <sup>59</sup>	Australia	RCT	Sub	Secondary	110	44	Overweight
Morgan <i>et al</i> , 2009 <sup>60</sup>	Australia	RCT	Obj	Secondary	65	36	Overweight
Patrick <i>et al</i> , 2011 <sup>61</sup>	USA	RCT	Sub	Secondary	441	44	Overweight
Petrella <i>et al</i> , 2017 <sup>62</sup>	Canada	RCT	Obj	Secondary	80	49	Overweight
Pritchard <i>et al</i> , 1997 <sup>63</sup>	Australia	RCT	Sub	Secondary	66	43	Overweight
Schröder <i>et al</i> , 2018 <sup>64</sup>	Spain	RCT	Sub	Secondary	6059	65	Overweight, MetS
Shin <i>et al</i> , 2017 <sup>65</sup>	Korea	RCT	Sub	Secondary	105	28	Overweight
Viestar <i>et al</i> , 2018 <sup>66</sup>	Netherlands	RCT	Sub	Secondary	314	47	General
Werkman <i>et al</i> , 2010 <sup>67</sup>	Netherlands	Cluster	Sub	Secondary	413	60	General
Wyke <i>et al</i> , 2019 <sup>68</sup>	Europe‡	RCT	Obj	Primary	1113	46	Overweight

\*Primary outcome indicates whether authors identified change in physical activity as a primary or secondary outcome of the study.

†Value for total sample; includes women in mixed-sex studies (ie, Gong *et al*<sup>48</sup>, 2015 (42% men), Werkman *et al*<sup>67</sup>, 2010 (85% men)).

‡England, Norway, Netherlands and Portugal.

Cluster, cluster randomised trial; MetS, metabolic syndrome; Obj, objective; PA, physical activity; RCT, randomised controlled trial; Sub, subjective.

about the included studies. Articles were published between 1997 and 2019 including a total sample size of 12 040 men. Ethnicity of participants were reported in only seven studies, of which five were predominantly white (>70%). Interventions primarily targeted overweight individuals (number of effect sizes ( $k$ )=14) and lasted on average 20 weeks (range 4–52 weeks). Intervention design varied between studies and often included multiple components with the majority including an aspect of face-to-face contact ( $k$ =19), education ( $k$ =24; that is, increasing knowledge or understanding), training ( $k$ =20; that is, imparting skills) and enablement ( $k$ =20; that is, increasing means/reducing barriers to increase capability or opportunity). Overall study attrition was 15% (range 5%–30%).

Online supplementary table S4 displays the study quality assessment for all studies. Overall study quality was mostly moderate ( $k$ =20; 83%), primarily due to participant self-referral (selection bias) and an inability to blind participants due to the nature of behavioural interventions. Studies rated as strong ( $k$ =4; 17%) were able to account for these issues by recruiting participants through a comprehensive list of the population (eg, clinical registry), which may not be feasible in community-based research and making efforts to blind assessors to participants' group allocation. The overall quality of evidence, assessed using GRADE,<sup>28 29</sup> was determined to be high for physical activity change and moderate for long-term physical activity change suggesting that we are very confident and moderately confident, respectively, that the true effect lies close to that of the estimate of the effect. Although there were concerns relating

to self-referral and an inability to blind participants in some studies, we did not downgrade the quality because we deemed the overall risk of bias to be very low. Long-term physical activity change was downgraded by one level for inconsistency based on considerable heterogeneity ( $I^2=80$ ) and relatively wide variance of point estimates across studies.

### Overall analysis—main findings

Intervention effects on physical activity are reported in figure 2. The estimated overall mean effect of physical activity interventions in men was small but significant ( $d=0.35$ ;  $SE=0.05$ ; 95% CI 0.26 to 0.45;  $p<0.001$ ). The effect size is consistent with a mean difference of 97 min of total physical activity per week or 980 steps per day between intervention and control participants. Review of the homogeneity statistic revealed a significant heterogeneous distribution ( $Q_w=72.32$ ,  $p<0.001$ ;  $I^2=68.20$ ). The one study removed procedure indicated that no individual study had a substantial impact on the overall effect size. Egger's regression test revealed that publication bias may be present ( $p<0.01$ ). No studies were added during the Trim and Fill procedure. Fail-safe N revealed that at least 876 unidentified studies with a mean effect of zero would be needed before the overall effect would no longer be statistically significant ( $p>0.05$ ).

Twelve studies<sup>45 47 48 50 51 53 56 59 60 66–68</sup> reported an additional follow-up measure at least 12 months postintervention (figure 3). The overall mean effect for long-term physical activity change was small but significant ( $d=0.32$ ;  $SE=0.09$ ; 95% CI 0.15 to

Table 2 Intervention characteristics

Primary source	Intervention					Types of behaviour change techniques (num; type)
	Delivery	Focus	Contact	Gender tailored	Duration (weeks)	
Aguiar <i>et al</i> , 2016 <sup>43</sup>	F2F, On	Com	Once	Yes	24	3 (Ed,T,En)
Andersen <i>et al</i> , 2012 <sup>44,45</sup>	F2F, Tel	PA	2–3/week	No	20	3 (Ed,T,En)
Ashton <i>et al</i> , 2017 <sup>46</sup>	F2F, On	Com	Weekly	Yes	12	3 (Ed,T,En)
Galvão <i>et al</i> , 2017 <sup>47</sup>	Tel	Com	Monthly	No	24	3 (Ed,Ev,En)
Gong <i>et al</i> , 2015 <sup>48</sup>	F2F, Tel	PA	Weekly	No	6	3 (Ed,T,Ev)
Gray <i>et al</i> , 2013 <sup>49</sup>	F2F	Com	Weekly	Yes	12	6 (Ed,I,T,Ev,M,En)
Groeneveld <i>et al</i> , 2011 <sup>50</sup>	F2F, Tel	Com	Monthly	No	24	3 (Ed,T,En)
Hunt <i>et al</i> , 2014 <sup>51</sup>	F2F	Com	Weekly	Yes	12	6 (Ed,I,T,Ev,M,En)
Livingston <i>et al</i> , 2015 <sup>52,53</sup>	F2F	PA	Biweekly	No	12	2 (Ed,T)
Maruyama <i>et al</i> , 2010 <sup>54</sup>	F2F, On, Tel	Com	Monthly	No	16	3 (Ed,T,En)
McGowan <i>et al</i> , 2013 <sup>55</sup>	Tel, Mail	PA	Once	No	4	2 (Ed,T)
Morgan <i>et al</i> , 2013 <sup>56</sup>	On, Mail	Com	Biweekly	Yes	12	3 (Ed,M,En)
Morgan <i>et al</i> , 2014 <sup>57</sup>	F2F	Com	Biweekly	Yes	7	4 (Ed,T,Ev,En)
Morgan <i>et al</i> , 2011a <sup>58</sup>	F2F	Com	2–3/month	Yes	12	4 (Ed,T,Ev,En)
Morgan <i>et al</i> , 2011b <sup>59</sup>	F2F, On	Com	2–3/month	Yes	12	5 (Ed,I,T,Ev,En)
Morgan <i>et al</i> , 2009 <sup>60</sup>	F2F, On	Com	2–3/month	Yes	12	3 (Ed,T,En)
Patrick <i>et al</i> , 2011 <sup>61</sup>	On	Com	Other	Yes	48	4 (Ed,T,Ev,En)
Petrella <i>et al</i> , 2017 <sup>62</sup>	F2F, On	Com	Weekly	Yes	12	5 (Ed,I,T,Ev,En)
Pritchard <i>et al</i> , 1997 <sup>63</sup>	F2F	PA	Bimonthly	No	48	2 (Ed,T)
Schröder <i>et al</i> , 2018 <sup>64</sup>	F2F, Tel	Com	3/month	No	52	3 (Ed,T,En)
Shin <i>et al</i> , 2017 <sup>65</sup>	F2F, On	Com	Monthly	No	12	3 (Ed,I,En)
Viestar <i>et al</i> , 2018 <sup>66</sup>	F2F, Tel	Com	Bi/monthly	No	24	3 (Ed,T,En)
Werkman <i>et al</i> , 2010 <sup>67</sup>	Mail, On	Com	Other	No	52	2 (Ed,En)
Wyke <i>et al</i> , 2019 <sup>68</sup>	F2F	Com	Weekly	Yes	12	6 (Ed,I,T,Ev,M,En)

Com, combined (eg, PA and diet); Ed, education; En, enablement; Ev, environmental; F2F, face-to-face; I, incentivisation; M, modelling; num, number; On, online; PA, physical activity; T, training; Tel, telephone.

0.48;  $p < 0.001$ ) and had a significant heterogeneous distribution ( $Q_w = 55.81$ ,  $p < 0.001$ ;  $I^2 = 80.29$ ). The one study removed procedure indicated that no individual study had a substantial impact on the overall effect size. Egger's regression test was non-significant ( $p > 0.05$ ). No studies were added during the Trim and Fill procedure and a Fail-safe N calculation indicated that 186 unidentified studies would be needed to nullify statistical significance ( $p > 0.05$ ).

## Moderator analyses

### Study characteristics

Mixed effects analysis produced significant between-moderator results for study characteristics on physical activity measurement,  $Q_b(1) = 9.30$ ,  $p \leq 0.01$ . Studies that employed objective measures of physical activity were found to have a larger effect size ( $d = 0.51$ ; 95% CI 0.37 to 0.65) than studies that used subjective measures of physical activity ( $d = 0.26$ ; 95% CI 0.17 to 0.35). Table 3 provides details of the analyses for study characteristics.




Figure 2 Forest plot of effect sizes representing effect on physical activity (baseline to postintervention).




Figure 3 Forest plot of effect sizes representing effect on long-term (ie,  $\geq 12$  months) physical activity change.

**Table 3** Physical activity interventions moderator analyses

	Effect size descriptive statistics					Null test	Heterogeneity statistics		
	k	d	SE	s <sup>2</sup>	95% CI		Q	T <sup>2</sup>	I <sup>2</sup>
Random effects model†	24	0.35	0.05	0.002	(0.26 to 0.45)	7.13***	72.32†	0.030	68.20
Study characteristics‡									
Study design							0.40		
RCT	21	0.37	0.05	0.003	(0.26 to 0.47)	6.80***		0.03	70.03
Cluster RCT	3	0.27	0.14	0.021	(-0.01 to 0.55)	1.87		0.05	62.55
Study quality							0.00		
Strong	4	0.36	0.13	0.002	(0.10 to 0.61)	2.75**		0.01	11.87
Moderate	20	0.35	0.05	0.003	(0.25 to 0.46)	6.51***		0.03	72.00
Measure							9.30**		
Subjective	14	0.26	0.05	0.002	(0.17 to 0.35)	5.66***		0.01	48.41
Objective	10	0.51	0.07	0.005	(0.37 to 0.65)	7.33***		0.01	29.81
PA outcome							0.33		
Primary	7	0.32	0.09	0.009	(0.14 to 0.50)	3.42***		0.03	63.97
Secondary	16	0.39	0.07	0.004	(0.26 to 0.51)	5.97***		0.04	68.91
Sample size							2.83		
n≤150	13	0.46	0.08	0.006	(0.30 to 0.61)	5.76***		0.02	29.30
n≥151	11	0.29	0.06	0.003	(0.18 to 0.41)	4.96***		0.03	77.89
Participant characteristics‡									
Mean age							0.20		
≤44 years	9	0.39	0.10	0.009	(0.20 to 0.58)	4.09***		0.00	0.00
≥45 years	15	0.34	0.06	0.003	(0.23 to 0.45)	5.88***		0.04	77.95
Population							0.78		
General	20	0.38	0.06	0.003	(0.27 to 0.49)	6.68***		0.04	71.28
Clinical	4	0.26	0.12	0.014	(0.03 to 0.49)	2.21*		0.02	47.50
Intervention characteristics‡									
Contact frequency							14.11***		
<1 weekly	14	0.22	0.05	0.002	(0.14 to 0.31)	4.98***		0.00	22.86
≥1 weekly	10	0.50	0.06	0.003	(0.39 to 0.62)	8.51***		0.02	47.60
Engagement							3.95		
High	3	0.68	0.17	0.028	(0.35 to 1.01)	4.04***		0.18	74.46
Moderate	10	0.34	0.07	0.010	(0.20 to 0.48)	4.79***		0.03	63.95
Low	6	0.31	0.10	0.005	(0.11 to 0.51)	3.04**		0.00	0.00
Gender-tailored							12.08***		
No	12	0.22	0.05	0.002	(0.12 to 0.31)	4.53***		0.00	21.27
Yes	12	0.47	0.06	0.003	(0.36 to 0.58)	8.55***		0.02	46.82
Duration							8.72**		
≤12 weeks	13	0.46	0.06	0.003	(0.35 to 0.57)	8.03***		0.03	53.08
≥13 weeks	11	0.23	0.05	0.003	(0.12 to 0.33)	4.26***		0.00	25.01
Theory							8.28**		
No	5	0.15	0.07	0.005	(0.01 to 0.30)	2.07*		0.00	0.01
Yes	19	0.40	0.05	0.002	(0.31 to 0.49)	8.84***		0.02	49.62
Behaviour change techniques							10.62***		
≤3 techniques	16	0.24	0.05	0.002	(0.15 to 0.34)	5.32***		0.01	26.98
≥4 techniques	8	0.51	0.07	0.004	(0.38 to 0.63)	7.72***		0.03	59.22

\*P≤0.05, \*\*P≤0.01, \*\*\*P≤0.001.

†Q<sub>w</sub> value used to determine heterogeneity.‡Q<sub>b</sub> value used to determine significant differences between moderators.d, effect size (Cohen's d); I<sup>2</sup>, total variance explained by moderator(s); k, number of effect sizes; s<sup>2</sup>, variance; SE, SE error; T<sup>2</sup>, between study variance in random effects model; Z, test of the null hypothesis.**Intervention characteristics**

Significant between-moderator results were present for contact frequency, Q<sub>b</sub>(1) = 14.11, p≤0.001, gender tailoring, Q<sub>b</sub>(1) = 12.08, p≤0.001, duration, Q<sub>b</sub>(1) = 8.72, p≤0.01, theory, Q<sub>b</sub>(1) = 8.28, p≤0.01 and number of types of behaviour change techniques, Q<sub>b</sub>(1) = 10.62, p≤0.001. Interventions that had between one or more weekly contacts produced a larger effect

size (d=0.50; 95% CI 0.39 to 0.62) than interventions that had less than 1 weekly contact (d=0.22; 95% CI 0.14 to 0.31). Interventions identified as gender-tailored had a larger effect size (d=0.47; 95% CI 0.36 to 0.58) than studies that were not gender-tailored (d=0.22; 95% CI 0.12 to 0.31). Those that were 12 weeks or less in duration produced a larger effect size (d=0.46; 95% CI 0.35 to 0.57) than interventions lasting 13

weeks or longer ( $d=0.23$ ; 95% CI 0.12 to 0.33). Interventions that identified one or more theory used the guide intervention design produced a larger effect size ( $d=0.40$ ; 95% CI 0.31 to 0.49) than interventions that did not use theory ( $d=0.15$ ; 95% CI 0.01 to 0.30). Interventions that used four or more types of behaviour change techniques had larger effect sizes ( $d=0.51$ ; 95% CI 0.38 to 0.63) than those that used three or less types of behaviour change techniques ( $d=0.24$ , 95% CI 0.15 to 0.34). All moderators had low between-study variance ( $T^2$ ) and explained moderate to large portions of subgroup variance ( $I^2$ ). Table 3 provides details of the analyses for intervention characteristics.

## DISCUSSION

This paper reports on the first meta-analysis of RCTs to synthesise the effects of behaviour change interventions to increase men's physical activity, sustained physical activity change (at 12 months) and to identify variations in outcomes due to potential moderating variables. Overall, interventions had a small but significant effect on increasing physical activity levels ( $d=0.35$ ) as well as postintervention sustainability of changes in physical activity ( $d=0.32$ ). The effect size is consistent with an increase of approximately 97 min of total physical activity per week or 980 steps per day between intervention and control participants. These effect sizes are larger than a meta-analysis investigating the effect of physical activity interventions for healthy adults ( $d=0.19$ ; 358 studies, 99 011 participants); however, the samples in that review were predominantly (74%) women.<sup>2</sup> Similar meta-analyses of interventions to promote physical activity among sedentary adults ( $d=0.31$ ; 11 studies; 3940 participants; 44% men)<sup>12</sup> and chronically ill (eg, hypertension, cancer, diabetes) adults ( $d=0.45$ ; 163 studies, 22 527 participants, 50% men)<sup>1</sup> are more in line with the present findings.

### The role of physical activity in men's health promotion

Our finding suggests that physical activity may be an important point-of-entry to encourage men's participation in behaviour change. As Connell<sup>69</sup> suggested, 'masculinity' is associated with action and doing, and physical activity clearly qualifies as an acceptable outlet and performance opportunity for men. Men are motivated to engage in activities which are perceived to be more 'masculine', such as sport and physical activity, as they may be associated with strength, friendly competition and mastery.<sup>70</sup> Particularly when the aim is to improve health, including a focus on physical activity may be more appealing and acceptable to men than providing support for 'dieting' or dietary modifications alone, as demonstrated in evidence from the UK and elsewhere.<sup>70-72</sup> Notably, a majority of the identified interventions in the present analysis had a primary focus on other health behaviours (eg, weight loss) but used physical activity as an adjunct intervention strategy. Further, many of the studies ( $k=19$ ; 79%) combined physical activity and other health behaviours (eg, diet), revealing opportunities for layering behaviour changes, in relative and relational ways that may engage and sustain men's participation. In this regard, physical activity may be viewed as a gateway that garners masculine capital, through which men may become more willing to address other health behaviours.<sup>20 73</sup>

### Participant characteristics and subpopulations

Participant characteristics within the included studies highlight some important trends and areas for future research. Although the target audience for many of the interventions reviewed here were overweight men, there are also likely to be gains in

tailoring physical activity programmes for other subgroups of men including those experiencing chronic illness. For example, prescription physical activity<sup>74</sup> and recreational football<sup>75</sup> have proven to be strong draws for men living with prostate cancer, as they have for weight loss<sup>51</sup> and physical activity interventions.<sup>68</sup> Resistance training has also been noted as a preferred modality of physical activity among men due to its perceived 'masculine' nature associated with strength but most importantly because it targets disease-related risk factors for men (ie, bone loss for men on androgen deprivation therapy).<sup>76</sup> Further, few studies provided sufficient detail of participants' sociocultural background, illuminating the need to formally evaluate the fit of men's physical activity health promotion programmes for marginalised subgroups. Men's health inequities, for example, may limit the access and involvement of subgroups including Indigenous men, those from culturally and linguistically diverse communities and men with low socioeconomic status. These subpopulations may benefit from culturally sensitive approaches to physical activity, with programmes designed to reduce structural barriers and address the resource-poor realities of these underserved end-users. It is especially poignant within these contexts that sustained programming, inclusive of longitudinal evaluations, are completed to ensure sustainability of the changes that are often initially garnered and gained immediately postintervention.

### Implications for intervention design

Our findings indicate that a variety of programme designs and approaches hold potential for positively influencing men's physical activity. For instance, using multiple appropriate behaviour change techniques and increased contact with participants (ie, at least weekly throughout the programme) were associated with significantly larger physical activity effects, suggesting that intensive interventions (ie, greater intervention dose and frequency) may be more effective. While brief or limited contact interventions may be appealing as a cost-effective option, they limit opportunities for men to interact and connect with similar others. There is evidence to suggest that men are drawn to programmes where they can connect with men similar to themselves, in a 'male-friendly' environment and engage in friendly banter and competition.<sup>20 72 77</sup> That shorter interventions (12 weeks or less) were more effective than longer interventions may reflect that the majority of reviewed studies ( $k=11$ ) were 12 weeks in duration. Although very brief interventions can produce acute changes in physical activity,<sup>78 79</sup> sustainable behaviour change likely requires some threshold of intervention intensity. Similarly, if the relative intervention intensity of longer duration interventions is too low, it may not be sufficient to elicit behaviour change. That most of these interventions ran for 12 weeks may bias or affirm that the optimal intervention duration for men lies somewhere in this range.

Although the majority of interventions in the included studies involved some type of face-to-face contact and multiple behaviour change techniques, it is noteworthy that interventions identified as gender-tailored were significantly more effective than those that were not ( $d=0.47$  vs 0.22). This encouraging evidence adds to a small but growing body of research that indicates that the mode of programme delivery, as well as the content, is an integral factor in successful programmes aimed at increasing men's physical activity.<sup>15 16</sup> While offering men-only interventions is a gender-tailored intervention in itself, the novel and diverse modes of programme delivery represented in these interventions point to the need to be responsive to the needs and



preferences of diverse groups of men to optimise intervention engagement and programme outcomes. Nevertheless, strategies used in the gender-tailored interventions reflect themes identified in successful efforts to promote men's health described by others and provide a useful direction for continuing efforts to promote men's health.<sup>18 19 80</sup> For example, gender-related strategies found to engage and retain men include the use of male-oriented language (eg, simple, straightforward messages/communications), images, humour and positive 'banter'; action oriented, strength-based approaches including realistic and manageable recommendations and providing men with flexible options that promote autonomy, self-reliance and mastery.

Interventions that were assessed using objective measures of physical activity were significantly more effective than interventions using subjective (self-report) measures. Self-reported behaviour has been observed to be both higher and lower than objective measures of physical activity.<sup>81</sup> Factors shown to predict discordances between measures include demographic characteristics such as education status<sup>82</sup> as well as differences in perceptions of what constitutes moderate or vigorous activity across demographic subgroups.<sup>83</sup> Interpreting subjective measures can be challenging.<sup>84 85</sup> For example, an intervention may affect how accurately an individual perceives and reports their physical activity and sedentary behaviour.<sup>84</sup>

### Sustaining long-term physical activity change

Though additional research is needed regarding long-term physical activity change, it is promising that the 12 of 24 studies that included a long-term (ie,  $\geq 12$  months) follow-up measure had a small intervention effect ( $d=0.32$ ). One study in the present analysis targeted overweight inactive male football fans and reported evidence of long-term behaviour maintenance of participants at 3.5-year follow-up.<sup>86</sup> These longitudinal findings of 213 men suggested that physical activity was significantly higher at 3.5 years than at baseline. This large-scale trial has informed the development of several subsequent gender-tailored interventions that also now include long-term follow-up in order to assess sustainability of physical activity change. For example, EuroFit,<sup>68</sup> which was delivered in four European countries, engaging 1113 men, includes a long-term follow-up of 12 months. Researchers must continue to evaluate the extent to which changes are sustained following intervention completion and consider strategies to promote long-term behaviour change.

### Strengths and limitations

We report two study strengths and three limitations. This meta-analysis is the first to examine the effects of behaviour change interventions in men and it builds on previous research identifying effective intervention strategies for engaging and retaining male participants. In addition, all 24 included studies had a randomised design, thus minimising bias.

The considerable heterogeneity across the studies, including the target population, the ways in which studies targeted men as participants, and the various modes of intervention delivery represent a limitation. We included both clinical (eg, patients with cancer) and healthy populations which may be viewed as a limitation even though physical activity has been effective in the setting of several different chronic illnesses<sup>1</sup> and healthy populations.

With regard to the assessment of study quality, we acknowledge that there is often poor agreement between tools and that tools may measure different constructs of study quality.<sup>25</sup> The EPHP is the recommended tool for assessing the quality of public health and

health promotion studies. Despite this, challenges with assessing community-based trials remain as the majority of studies were rated down for participant self-referral and a lack of participant blinding. For almost any real-world behavioural intervention, a degree of volition and motivation to attend is required, both overall and session-by-session. Further, usually it is not possible to blind participants, or those delivering or assessing the intervention to group allocation, in these type of interventions (ie, participants know they are exercising). Considered collectively, these assessment tools include assumptions which favour a biomedical approach in which internal validity, as an outcome of tightly controlled trials, is prioritised over external validity and thus raises challenges for real-world implementation.<sup>87</sup> What is not reflected within such assessment is the value of pragmatic trials in implementing behaviour change within the environments in which they will be used.<sup>87 88</sup>

We also acknowledge that potentially relevant studies conducted on mixed-sex samples were not included because they failed to disaggregate their findings by sex—despite calls for the need to do so.<sup>9</sup> Thus, the majority of studies included had male participants only.

### CONCLUSION

This meta-analysis suggests that men can make small, but potentially important, changes in their physical activity. Specifically, gender-tailored interventions, which include a core focus on physical activity, may help attract, engage and retain men to health behaviour interventions. This includes those with a primary focus on other behaviours, which is important for the improvement of individual and public health, in addition to physical activity. Scope clearly exists for researchers to better understand what types of interventions work for different men and why. Sustained improvement in physical activity is a public health holy grail and our study captures the current knowledge about men in that domain.

#### What is already known

- ▶ Men are under-represented in health promotion programmes.
- ▶ Physical activity has been identified as a draw facilitating men's engagement with health promotion initiatives and programmes and/or injury and illness management.
- ▶ Growing interest in men's health promotion has led to the development of interventions which include a focus on physical activity.
- ▶ Two systematic reviews have been conducted on physical activity interventions in men; yet the effects of randomised controlled trial (RCT) physical activity interventions are poorly understood.

#### What are the new findings

- ▶ Behaviour change interventions have a small but significant positive effect on men's physical activity.
- ▶ Interventions that (i) are based on a theoretical framework, (ii) are tailored to men's values and interests, (iii) include regular group contact and (iv) employ multiple behaviour change strategies appear most effective.
- ▶ There is some evidence that physical activity is sustained at long-term follow-up but more longitudinal research is needed.

**Twitter** Paul Sharp @paulgsharp

**Contributors** PS, JCS, JLB, JLO, KH, MVD and CMC helped conceive the project and PS and JCS were responsible for project management. PS, MVD and CMC



determined the search strategy and oversaw the screening of articles. PS and JCS were responsible for data extraction, analysis and interpretation. PS and CMC undertook quality assessment. All authors participated in the study design, drafting of the manuscript and critical revisions. All authors read and approved the final manuscript.

**Funding** PS is supported by an Australian Government Research Training Programme Scholarship.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## ORCID iD

Paul Sharp <http://orcid.org/0000-0001-5616-3181>

## REFERENCES

- Conn VS, Hafidahl AR, Brown SA, *et al.* Meta-Analysis of patient education interventions to increase physical activity among chronically ill adults. *Patient Educ Couns* 2008;70:157–72.
- Conn VS, Hafidahl AR, Mehr DR. Interventions to increase physical activity among healthy adults: meta-analysis of outcomes. *Am J Public Health* 2011;101:751–8.
- Evans J, Frank B, Oliffe JL, *et al.* Health, illness, men and masculinities (HIMM): a theoretical framework for understanding men and their health. *J Mens health* 2011;8:7–15.
- Wang H, Naghavi M, Allen C, *et al.* Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the global burden of disease study 2015. *The Lancet* 2016;388:1459–544.
- Courtenay WH. Constructions of masculinity and their influence on men's well-being: a theory of gender and health. *Soc Sci Med* 2000;50:1385–401.
- Galdas PM, Cheater F, Marshall P. Men and health help-seeking behaviour: literature review. *J Adv Nurs* 2005;49:616–23.
- Levant RF, Wimer DJ. Masculinity constructs as protective buffers and risk factors for men's health. *Am J Mens Health* 2014;8:110–20.
- Yousaf O, Grunfeld EA, Hunter MS. A systematic review of the factors associated with delays in medical and psychological help-seeking among men. *Health Psychol Rev* 2015;9:264–76.
- Waters LA, Galichet B, Owen N, *et al.* Who participates in physical activity intervention trials? *J Phys Act Health* 2011;8:85–103.
- Hallal PC, Andersen LB, Bull FC, *et al.* Global physical activity levels: surveillance progress, pitfalls, and prospects. *The Lancet* 2012;380:247–57.
- Warburton DER, Bredin SSD. Health benefits of physical activity: a systematic review of current systematic reviews. *Curr Opin Cardiol* 2017;32:541–56.
- Foster C, Hillsdon M, Thorogood M, *et al.* Interventions for promoting physical activity. *Cochrane Database Syst Rev* 2005;18.
- Foster C, Richards J, Thorogood M, *et al.* Remote and web 2.0 interventions for promoting physical activity. *Cochrane Database of Systematic Reviews* 2013.
- Richards J, Hillsdon M, Thorogood M, *et al.* Face-to-face interventions for promoting physical activity. *Cochrane Database of Systematic Reviews* 2013.
- Bottoff JL, Seaton CL, Johnson ST, *et al.* An updated review of interventions that include promotion of physical activity for adult men. *Sports Med* 2015;45:775–800.
- George ES, Kolt GS, Duncan MJ, *et al.* A review of the effectiveness of physical activity interventions for adult males. *Sports Medicine* 2012;42:281–300.
- Hunt K, Gray CM, Maclean A, *et al.* Do weight management programmes delivered at professional football clubs attract and engage high risk men? A mixed-methods study. *Public Health* 2014;14:50.
- Morgan PJ, Warren JM, Lubans DR, *et al.* Engaging men in weight loss: experiences of men who participated in the male only SHED-IT pilot study. *Obes Res Clin Pract* 2011;5:e239–48.
- Robertson LM, Douglas F, Ludbrook A, *et al.* What works with men? A systematic review of health promoting interventions targeting men. *BMC Health Serv Res* 2008;8:141.
- Sharp P, Bottoff JL, Hunt K, *et al.* Men's Perspectives of a Gender-Sensitized Health Promotion Program Targeting Healthy Eating, Active Living, and Social Connectedness. *American Journal of Men's Health* 2018;0:1557988318799159.
- Moher D, Liberati A, Tetzlaff J. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.
- Bozzo A, Ball K, Evaniew N, *et al.* Retractions in cancer research: a systematic survey. *Research Integrity and Peer Review* 2017;2:5.
- Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implementation Science* 2011;6:42.
- Effective Public Health Practice Project. Quality assessment tool for quantitative studies Hamilton, on: effective public health practice project, 1998. Available: <https://merst.ca/ephpp/>
- Armijo-Olivo S, Stiles CR, Hagen NA, *et al.* Assessment of study quality for systematic reviews: a comparison of the Cochrane collaboration risk of bias tool and the effective public health practice project quality assessment tool: methodological research. *J Eval Clin Pract* 2012;18:12–18.
- Thomas BH, Ciliska D, Dobbins M, *et al.* A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews Evid Based Nurs* 2004;1:176–84.
- Jackson N, Waters E, Guidelines for Systematic Reviews in Health Promotion and Public Health Taskforce. Criteria for the systematic review of health promotion and public health interventions. *Health Promot Int* 2005;20:367–74.
- Balshem H, Helfand M, Schünemann HJ, *et al.* Grade guidelines: 3. rating the quality of evidence. *J Clin Epidemiol* 2011;64:401–6.
- GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ* 2004;328:1490.
- Cohen J. *Statistical power analysis for the behavioral sciences*, 1988.
- Green S, Higgins J. *Cochrane Handbook for systematic reviews of interventions: cochrane training*, 2005.
- Borenstein M, Hedges LV, Higgins JP, *et al.* *Introduction to meta-analysis*. John Wiley & Sons, 2011.
- Rao JNK, Scott AJ. A simple method for the analysis of clustered binary data. *Biometrics* 1992;48:577–85.
- Heck RH, Tabata L, Thomas SL. *Multilevel and longitudinal modeling with IBM SPSS*. Routledge, 2013.
- Kianoush F, Masoomi K. Application REML model and determining cut off of ICC by multi-level model based on Markov chains simulation in health. *Indian Journal of Fundamental and Applied Life Sciences* 2015;5:1432–48.
- Orron G, Kinmonth A-L, Sanderson S, *et al.* Effectiveness of physical activity promotion based in primary care: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2012;344:e1389.
- Elley CR, Kersner N, Arroll B. Effectiveness of counselling patients on physical activity in general practice: cluster randomised controlled trial. *BMJ* 2003;326:793.
- Egger M, Smith GD, Schneider M, *et al.* Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- Duval S, Tweedie R. Trim and fill: a simple Funnel-Plot-Based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56:455–63.
- Rosenthal R. *Meta-Analytic procedures for social research*. Sage, 1991.
- Davies CA, Spence JC, Vandelandotte C, *et al.* Meta-Analysis of internet-delivered interventions to increase physical activity levels. *Int J Behav Nutr Phys Act* 2012;9:52.
- Greaves CJ, Sheppard KE, Abraham C, *et al.* Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC Public Health* 2011;11:119.
- Aguiar EJ, Morgan PJ, Collins CE, *et al.* Efficacy of the type 2 diabetes prevention using lifestyle education program RCT. *Am J Prev Med* 2016;50:353–64.
- Andersen E, Burton NW, Andersen SA. Physical activity levels six months after a randomised controlled physical activity intervention for Pakistani immigrant men living in Norway. *Int J Behav Nutr Phys Act* 2012;9:47.
- Andersen E, Høstmark AT, Holme I, *et al.* Intervention effects on physical activity and insulin levels in men of Pakistani origin living in Oslo: a randomised controlled trial. *J Immigr Minor Health* 2013;15:101–10.
- Ashton LM, Morgan PJ, Hutchesson MJ, *et al.* Feasibility and preliminary efficacy of the 'HEYMAN' healthy lifestyle program for young men: a pilot randomised controlled trial. *Nutr J* 2017;16:2.
- Galvão DA, Newton RU, Giris A, *et al.* Randomized controlled trial of a peer led multimodal intervention for men with prostate cancer to increase exercise participation. *Psychooncology* 2018;27:199–207.
- Gong J, Chen X, Li S. Efficacy of a community-based physical activity program KM2H2 for stroke and heart attack prevention among senior hypertensive patients: a cluster randomized controlled phase-II trial. *PLoS One* 2015;10:e0139442.
- Gray CM, Hunt K, Mutrie N, *et al.* Weight management for overweight and obese men delivered through professional football clubs: a pilot randomized trial. *Int J Behav Nutr Phys Act* 2013;10:121.
- Groeneveld IF, Proper KI, van der Beek AJ, *et al.* Short and long term effects of a lifestyle intervention for construction workers at risk for cardiovascular disease: a randomized controlled trial. *BMC Public Health* 2011;11:836.
- Hunt K, Wyke S, Gray CM, *et al.* A gender-sensitized weight loss and healthy living programme for overweight and obese men delivered by Scottish premier league football clubs (FFIT): a pragmatic randomised controlled trial. *Lancet* 2014;383:1211–21.
- Livingston PM, Craike MJ, Salmon J, *et al.* Effects of a clinician referral and exercise program for men who have completed active treatment for prostate cancer: a multicenter cluster randomized controlled trial (engage). *Cancer* 2015;121:2646–54.
- Gaskin CJ, Craike M, Mohebbi M, *et al.* A clinician referral and 12-week exercise training program for men with prostate cancer: outcomes to 12 months of the engage cluster randomized controlled trial. *J Phys Act Health* 2017;14:353–9.
- Maruyama C, Kimura M, Okumura H, *et al.* Effect of a worksite-based intervention program on metabolic parameters in middle-aged male white-collar workers: a randomized controlled trial. *Prev Med* 2010;51:11–17.

- 55 McGowan EL, North S, Courneya KS. Randomized controlled trial of a behavior change intervention to increase physical activity and quality of life in prostate cancer survivors. *Ann Behav Med* 2013;46:382–93.
- 56 Morgan PJ, Callister R, Collins CE, *et al.* The SHED-IT community trial: a randomized controlled trial of internet- and paper-based weight loss programs tailored for overweight and obese men. *Ann Behav Med* 2013;45:139–52.
- 57 Morgan PJ, Collins CE, Plotnikoff RC, *et al.* The 'Healthy Dads, Healthy Kids' community randomized controlled trial: A community-based healthy lifestyle program for fathers and their children. *Prev Med* 2014;61:90–9.
- 58 Morgan PJ, Lubans DR, Callister R, *et al.* The 'Healthy Dads, Healthy Kids' randomized controlled trial: efficacy of a healthy lifestyle program for overweight fathers and their children. *Int J Obes* 2011;35:436–47.
- 59 Morgan PJ, Collins CE, Plotnikoff RC, *et al.* Efficacy of a workplace-based weight loss program for overweight male shift workers: the workplace power (preventing obesity without eating like a rabbit) randomized controlled trial. *Prev Med* 2011;52:317–25.
- 60 Morgan PJ, Lubans DR, Collins CE, *et al.* The SHED-IT randomized controlled trial: evaluation of an Internet-based weight-loss program for men. *Obesity* 2009;17:2025–32.
- 61 Patrick K, Calfas KJ, Norman GJ, *et al.* Outcomes of a 12-month web-based intervention for overweight and obese men. *Ann Behav Med* 2011;42:391–401.
- 62 Petrella RJ, Gill DP, Zou G, *et al.* Hockey fans in training: a pilot pragmatic randomized controlled trial. *Med Sci Sports Exerc* 2017;49:2506–16.
- 63 Pritchard JE, Nowson CA, Wark JD. A worksite program for overweight middle-aged men achieves lesser weight loss with exercise than with dietary change. *J Am Diet Assoc* 1997;97:37–42.
- 64 Schröder H, Cárdenas-Fuentes G, Martínez-González MA, *et al.* Effectiveness of the physical activity intervention program in the PREDIMED-Plus study: a randomized controlled trial. *Int J Behav Nutr Phys Act* 2018;15:110.
- 65 Shin DW, Yun JM, Shin J-H, *et al.* Enhancing physical activity and reducing obesity through smartcare and financial incentives: a pilot randomized trial. *Obesity* 2017;25:302–10.
- 66 Viester L, Verhagen EALM, Bongers PM, *et al.* Effectiveness of a worksite intervention for male construction workers on dietary and physical activity behaviors, body mass index, and health outcomes: results of a randomized controlled trial. *Am J Health Promot* 2018;32:795–805.
- 67 Werkman A, Hulshof PJM, Stafleu A, *et al.* Effect of an individually tailored one-year energy balance programme on body weight, body composition and lifestyle in recent retirees: a cluster randomised controlled trial. *BMC Public Health* 2010;10:110.
- 68 Wyke S, Bunn C, Andersen E, *et al.* The effect of a programme to improve men's sedentary time and physical activity: The European Fans in Training (EuroFIT) randomised controlled trial. *PLoS Med* 2019;16:e1002736.
- 69 Connell RW. *Masculinities polity*, 2005.
- 70 Hunt K, McCann C, Gray CM, *et al.* "You've got to walk before you run": positive evaluations of a walking program as part of a gender-sensitized, weight-management program delivered to men through professional football clubs. *Health Psychol* 2013;32:57–65.
- 71 Archibald D, Douglas F, Hoddinott P, *et al.* A qualitative evidence synthesis on the management of male obesity. *BMJ Open* 2015;5:e008372.
- 72 Hunt K, Wyke S, Bunn C, *et al.* Scale-Up and scale-out of a gender-sensitized weight management and healthy living program delivered to overweight men via professional sports clubs: the wider implementation of football fans in training (FFIT). *Int J Environ Res Public Health* 2020;17:584.
- 73 Gough B. The psychology of men's health: maximizing masculine capital. *Health Psychol* 2013;32:1–4.
- 74 Cormie P, Oliffe JL, Wootten AC, *et al.* Improving psychosocial health in men with prostate cancer through an intervention that reinforces masculine values - exercise. *Psychooncology* 2016;25:232–5.
- 75 Bjerre E, Bruun DM, Tolver A, *et al.* Effectiveness of community-based football compared to usual care in men with prostate cancer: protocol for a randomised, controlled, parallel group, multicenter superiority trial (the Fc prostate community trial). *BMC Cancer* 2016;16:767.
- 76 Finlay A, Wittert G, Short CE. A systematic review of physical activity-based behaviour change interventions reaching men with prostate cancer. *J Cancer Surviv* 2018;12:571–91.
- 77 Bunn C, Wyke S, Gray CM, *et al.* 'Coz football is what we all have': masculinities, practice, performance and effectiveness in a gender-sensitized weight-loss and healthy living programme for men. *Social Health Illn* 2016;38:812–28.
- 78 GC V, Wilson ECF, Suhrcke M, *et al.* Are brief interventions to increase physical activity cost-effective? A systematic review. *Br J Sports Med* 2016;50:408–17.
- 79 National Institute for Clinical Excellence. *Four commonly used methods to increase physical activity: brief interventions in primary care, exercise referral schemes, pedometers and community-based exercise programmes for walking and cycling*. Great Britain: NICE, 2006.
- 80 Robertson S, Witty K, Zwolinsky S, *et al.* Men's health promotion interventions: what have we learned from previous programmes. *Community Pract* 2013;86:38–42.
- 81 Prince SA, Adamo KB, Hamel M, *et al.* A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5:56.
- 82 Gorzelitz J, Peppard PE, Malecki K, *et al.* Predictors of discordance in self-report versus device-measured physical activity measurement. *Ann Epidemiol* 2018;28:427–31.
- 83 Rebustini F, Balbinotti MAA, REdl F-R, *et al.* Sport psychometry, participants and invariance: a critical review. *Journal of Physical Education* 2016;27.
- 84 Kelly P, Fitzsimons C, Baker G. Should we reframe how we think about physical activity and sedentary behaviour measurement? validity and reliability reconsidered. *Int J Behav Nutr Phys Act* 2016;13:32.
- 85 Steene-Johannessen J, Anderssen SA, van der Ploeg HP, *et al.* Are self-report measures able to define individuals as physically active or inactive? *Med Sci Sports Exerc* 2016;48:235–44.
- 86 Gray CM, Wyke S, Zhang R, *et al.* Long-Term weight loss following a randomised controlled trial of a weight management programme for men delivered through professional football clubs: the football fans in training follow-up study. *Public Health Res* 2018;6:1–114.
- 87 Ammerman A, Smith TW, Calancie L. Practice-Based evidence in public health: improving reach, relevance, and results. *Annu Rev Public Health* 2014;35:47–63.
- 88 Glasgow RE, Vinson C, Chambers D, *et al.* National Institutes of health approaches to dissemination and implementation science: current and future directions. *Am J Public Health* 2012;102:1274–81.



## PRISMA 2009 Checklist

Section/topic	#	Checklist item
<b>TITLE</b>		
Title	1	Identify the report as a systematic review, meta-analysis, or both.
<b>ABSTRACT</b>		
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; participants, and interventions; study appraisal and synthesis methods; results; limitations; implications of key findings; systematic review registration number.
<b>INTRODUCTION</b>		
Rationale	3	Describe the rationale for the review in the context of what is already known.
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, outcomes, and study design (PICOS).
<b>METHODS</b>		
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and registration information including registration number.
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., language, publication status) used as criteria for eligibility, giving rationale.
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to request additional studies) in the search and date last searched.
Search	8	Present full electronic search strategy for at least one database, including any limits used, and how they were repeated.
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and included in the meta-analysis).
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate), and how data were confirmed or confirmed with investigators.
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any exclusions or simplifications made.
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specific methods for assessing risk of bias), and how this information is to be used in any data synthesis.
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of conflict (e.g., $I^2$ ) for each meta-analysis.
<b>RESULTS</b>		
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, reporting within studies).
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), which were pre-specified.

Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, design, location, etc.), and provide the citations.
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessments (e.g., risk of bias summary, risk of bias graph).
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary of effects and (b) effect estimates and confidence intervals, ideally with a forest plot.
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression).
<b>DISCUSSION</b>		
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome and important key groups (e.g., healthcare providers, users, and policy makers).
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., identified research, reporting bias).
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for practice.
<b>FUNDING</b>		
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of information, access to participants, etc.).

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097.doi:10.1371/journal.pmed1000097

For more information, visit: [www.prisma-statement.org](http://www.prisma-statement.org).



## Supplementary 2 (S2)

### Medline (Ovid SP) (1946-)

	Search Terms
1	Randomized Controlled Trial/
2	((randomi#ed or experimental) adj3 trial).ab,ti,kw or randomi#ed controlled trial.pt
3	Motor Activity/ or exp sports/ or Exercise/ or Circuit-based Exercise/ or High-intensity Interval Training/ or Muscle Stretching Exercises/ or Physical Conditioning, Human/ or Plyometric Exercise/ or Resistance Training/ or Exercise Therapy/ or Healthy Lifestyle/ or Health Promotion/
4	((((trial or program* or intervention?) adj4 (PA or "physical activit*" or exercis* or walk* or run* or jog* or sport* or swim* or cycl* or soccer or football or rugby or hockey or cricket or basketball or tennis or volleyball or golf or yoga)) or active living or active commuting or exercise therapy or healthy lifestyle or lifestyle change or lifestyle behavio?r or behavio?r change or health promotion or health behavio?r).ab,ti,kw
5	Program Evaluation/ or Self Report/ or "Surveys and Questionnaires"/ or Fitness Trackers/
6	("outcome assessment" or evaluation? or self-report* or survey? or questionnaire? or accelerometer? or pedometer? or self-monitoring or telephone or web-based or fitness tracker? or activity tracker? or ((physical activity or exercise) adj3 (index* or level?))).ab,ti,kw
7	Male/
8	(men or male?).ab,ti,kw
9	1 or 2
10	3 or 4
11	5 or 6
12	7 or 8
13	9 and 10 and 11 and 12
14	13 limited to English and Abstract

## Supplementary 3 (S3)

**Table S3. Brief overview of intervention and intervention focus**

Primary source	Intervention overview	Theoretical orientation	Intervention engagement	Measurement instrument	Measure
Aguiar <i>et al</i> , 2016(43)	<ul style="list-style-type: none"> <li>- The Prevention Using Lifestyle Education (PULSE) program consisted of:               <ul style="list-style-type: none"> <li>o Intervention content from SHED-IT in Morgan <i>et al.</i> (2013)</li> <li>o Diabetes Prevention Handbook for men and PULSE Exercise Support Book for Men, including prescribed home-based exercise program, logbook, and resistance band device</li> </ul> </li> </ul>	SCT	LOW 13% of men met log book compliance	Pedometer	Steps/day
Andersen <i>et al</i> , 2012(44, 45)	<ul style="list-style-type: none"> <li>- Structured group exercise sessions led by exercise physiologist twice a week</li> <li>- Two group lectures; individual counselling session; written material and a phone call</li> </ul>	SCT	LOW 60% attendance of exercise portion 90% attended group lectures All participants completed individual counselling and phone call.	Actigraph Accelerometer	Counts per minute per day
Ashton <i>et al</i> , 2017(46)	<ul style="list-style-type: none"> <li>- The HEYMAN program consisted of:               <ul style="list-style-type: none"> <li>o Responsive website</li> <li>o Wearable physical activity tracker and resistance band for home-based strength training with associated app/website.</li> <li>o One-hour weekly face-to-face exercise sessions (11x group based and 1x individual) focusing on aerobic and strength exercise</li> <li>o Personalised food and nutrient report</li> <li>o Private Facebook discussion group</li> </ul> </li> </ul>	SCT SDT	MODERATE 100% visited website at least once, 63% visited weekly, 58% used app daily, 91% attended one 60-min face-to-face session	Pedometer	Steps/day
Galvao <i>et al</i> , 2017(47)	<ul style="list-style-type: none"> <li>- Self-management materials and monthly telephone group (~6-8 participants) peer support for 6 months. 1/6 sessions focused on physical activity and peers discussed exercise at each session to check how men's physical activity goals were progressing</li> </ul>	Nil	MODERATE Mean attendance 4/6 (67%) sessions, 28% accessed website at least once	Godin Leisure-Time Exercise Questionnaire	Total exercise min/week

	<ul style="list-style-type: none"> <li>- Participants provided with elastic exercise device and heart rate monitor</li> <li>- Online material also available and men received feedback on unmet supportive care needs, distress score, waist circumference, and exercise levels</li> </ul>				
Gong <i>et al</i> , 2015(48)	<ul style="list-style-type: none"> <li>- The KM2H program consisted of:               <ul style="list-style-type: none"> <li>o Two 45–60-minute lectures (i.e., “Secrets for Blood Pressure Control through Exercise”; “Individualised Physical Activity Counselling”)</li> <li>o Two 10–20-minute telephone counselling on obstacles to engaging in physical activity</li> <li>o Two group sessions (~8–10 participants) provide opportunities to share skills, experiences, and build social support</li> <li>o Two booster sessions post-intervention</li> </ul> </li> </ul>	TTM MPM SCapT	unknown	Self-report Questionnaire (non-validated survey)	Levels of physical activity
Gray <i>et al</i> , 2013(49)	<ul style="list-style-type: none"> <li>- Football Fans in Training (FFIT) included twelve weekly 90-minute group (~15 men) sessions at professional football stadia.               <ul style="list-style-type: none"> <li>o Classroom-based education on topics related to weight management, healthy eating, alcohol and increasing daily physical activity</li> <li>o Coach-led physical activity sessions including training in aerobic, strength and flexibility exercises</li> </ul> </li> <li>- Pedometers to support self-monitoring as part of incremental pedometer-based walking program</li> </ul>	SCT CT	MODERATE 76% attended at least 80% of the sessions	International Physical Activity Questionnaire (Short Form)	Total activity (MET min/week)
Groeneveld <i>et al</i> , 2011(50)	<ul style="list-style-type: none"> <li>- Health under Construction consisted of:               <ul style="list-style-type: none"> <li>o Three 45-60-minute face-to-face sessions</li> <li>o Four 15-30-minute one-on-one telephone counselling sessions, using motivational interviewing delivered by occupational physician</li> <li>o Participant’s risk profile, personal determinants, and barriers for behaviour change were discussed, and goals set. Participants chose to aim at either diet and PA, or smoking</li> </ul> </li> </ul>	Nil	unknown	Short Questionnaire to Assess Health enhancing PA (SQUASH)	Leisure time physical activity and spot-related expenditure (MET min/week)
Hunt <i>et al</i> , 2014(51)	<ul style="list-style-type: none"> <li>- As for FFIT in Gray <i>et al</i>. (2013)</li> </ul>	SCT CT	MODERATE 79% attended at least 6/12 sessions	International Physical Activity	Total activity (MET min/week)

				Questionnaire (Short Form)	
Livingston <i>et al</i> , 2015(52, 53)	<ul style="list-style-type: none"> <li>- The ENGAGE intervention was a clinician-referred 12-week community exercise program consisting of:               <ul style="list-style-type: none"> <li>o Two 50-minute supervised gym sessions and one home-based session per week based on exercise guidelines for cancer survivors developed by the American College of Sports Medicine and Exercise and Sport Science Australia</li> </ul> </li> </ul>	SCT	HIGH 85% attended at least 18/24 gym session, of those that completed diary (74%), 81% completed 9-12 home-based weekly session	Godin Leisure-Time Exercise Questionnaire	MVPA min/week
Maruyama <i>et al</i> , 2010(54)	<ul style="list-style-type: none"> <li>- The LiSM program included:               <ul style="list-style-type: none"> <li>o Individualised assessment and collaborative goal setting sessions based on food group intake and physical activity</li> <li>o Two individual counselling sessions with a registered dietitian and physical trainer</li> <li>o Monthly website advice designed to promote healthy dietary habits and physical activity</li> <li>o Monthly individual contact with a dietitian and a physical trainer. The fourth counselling session, at end of the third month, conducted through the website</li> <li>o Personal and interactive website to self-monitor weight, food intake and physical activity, and discuss awareness of lifestyles for self-monitoring throughout the intervention period</li> <li>o Pedometer linked to website. Data automatically presented in figures on their individual website pages</li> </ul> </li> </ul>	Nil	LOW Adherence to self-reporting was low (75% of intervention, 50% control returning usable records)	Pedometer	Steps/day
McGowan <i>et al</i> , 2013(55)	<ul style="list-style-type: none"> <li>- The PROMOTE intervention provided all participants with physical activity Guidelines for Americans (2008)</li> <li>- Self-administered implementation group:               <ul style="list-style-type: none"> <li>o Recorded current levels of physical activity, set goals and made detailed plans to meet the physical activity guidelines, or increase physical activity by 60+ min/week if already</li> </ul> </li> </ul>	TPB	LOW 46% saying they did not really complete it, 23% reporting completing some of it, 17% most of it, and 14% all of it	Godin Leisure-Time Exercise Questionnaire	Total activity (min/week)



	<p>meeting guidelines. List personal physical activity barriers, and strategies to overcome barriers</p> <ul style="list-style-type: none"> <li>- The telephone-assisted implementation group: <ul style="list-style-type: none"> <li>o Same as above plus a 12-minute telephone call to assist with goal setting and planning</li> </ul> </li> </ul>				
Morgan <i>et al</i> , 2013(56)	<ul style="list-style-type: none"> <li>- Self-help, Exercise, Diet and Information Technology (SHED-IT) weight loss program for men consisted of a resource package including DVD, gender-tailored handbook and a pedometer to monitor step count <ul style="list-style-type: none"> <li>o Two of nine evidence-based weight loss messages focused on physical activity ('Every step counts', 'Reduce sitting time'. One of the three SMART goals that men were asked to set each month was related to physical activity</li> </ul> </li> <li>- Online version included exercise diary and some exercise-related feedback</li> </ul>	SCT	unknown	Pedometer	Steps/day
Morgan <i>et al</i> , 2014(57)	<ul style="list-style-type: none"> <li>- Health Dads, Healthy Kids (HDHK) consisted of seven weekly group sessions (90 minutes each): <ul style="list-style-type: none"> <li>o Four sessions for fathers only, and three practical sessions for fathers and children, which included co-physical activity.</li> <li>o Weight loss component of HDHK intervention was adapted from SHED-IT program</li> </ul> </li> </ul>	SCT FST	MODERATE Mean attendance rate for 7 sessions was 71%	Pedometer	Steps/day
Morgan <i>et al</i> , 2011a(58)	<ul style="list-style-type: none"> <li>- As for HDHK in Morgan <i>et al</i> (2014)</li> </ul>	SCT FST	HIGH Participants attended 81% of the sessions	Pedometer	Mean steps/day
Morgan <i>et al</i> , 2011b(59)	<ul style="list-style-type: none"> <li>- Workplace POWER (Preventing Obesity Without Eating like a Rabbit) consisted of: <ul style="list-style-type: none"> <li>o One face-to-face information session (75 min) covering energy balance, shift work challenges to diet and physical activity, weight loss tips, and behaviour change strategies (self-monitoring, goal setting, social support)</li> <li>o Group-based financial incentives (\$AU50 voucher per person for sport store for group with highest mean weight loss at 1 month and end of program)</li> <li>o Online component (direction to publicly accessible, free weight loss website)</li> </ul> </li> </ul>	SCT	LOW Average of 24 exercise entries on website across 14 weeks. 28% complied with web-based components	Godin Leisure-Time Exercise Questionnaire	Total MET minutes/week

	<ul style="list-style-type: none"> <li>Program booklet, weight loss handbook, and pedometer</li> </ul>				
Morgan <i>et al</i> , 2009(60)	<ul style="list-style-type: none"> <li>As for SHED-IT in Morgan <i>et al</i> (2013)</li> </ul>	SCT	LOW Average of 23 exercise entries on website throughout intervention. 41% complied with web-based tasks	Pedometer	Mean steps/day
Patrick <i>et al</i> , 2011(61)	<ul style="list-style-type: none"> <li>Internet-based intervention designed to promote weight loss through goals for diet and physical activity including increasing steps per day, and strength training consisting of:               <ul style="list-style-type: none"> <li>Computerised assessment to allow tailoring of messages</li> <li>Weekly web-based learning presented in short sessions using “business-like” language and graphics</li> <li>Individualised feedback on progress</li> <li>Pedometers to self-monitor daily step count and recording of minutes of other physical activity</li> </ul> </li> </ul>	SCT	MODERATE Average of 23.4 weekly logins	International Physical Activity Questionnaire (Long Form)	Total walking (min/day)
Petrella <i>et al</i> , 2017(62)	<ul style="list-style-type: none"> <li>As for FFIT in Gray <i>et al</i> [20] and Hunt <i>et al</i> [22], except ice-hockey setting</li> </ul>	CT SCT SRT	HIGH Mean session attendance of 82%	Pedometer	Mean steps/day
Pritchard <i>et al</i> , 1997(63)	<ul style="list-style-type: none"> <li>Two intervention arms including weight loss through diet or through physical activity               <ul style="list-style-type: none"> <li>Self-selected aerobic leisure exercise regimen, minimum 3 x weekly</li> </ul> </li> </ul>	Nil	MODERATE >70% attendance at bimonthly presentations	Activity Diary	Index of activity
Schröder <i>et al</i> , 2018(64)	<ul style="list-style-type: none"> <li>The PREDIMED-Plus program targeted goal setting, action planning, feedback, informational materials, motivation, and self-monitoring during:               <ul style="list-style-type: none"> <li>Twelve individual (60-minute) sessions</li> <li>Twelve telephone calls</li> <li>Three one-hour group sessions. Tailored physical activity goals and an action plan</li> </ul> </li> <li>Participants are provided with a pedometer and a physical activity diary for self-monitoring</li> </ul>	Nil	unknown	Regicor Short Physical Activity Questionnaire	MET min/day
Shin <i>et al</i> , 2017(65)	<ul style="list-style-type: none"> <li>The EPAROSFI program included:               <ul style="list-style-type: none"> <li>Standardised education materials</li> </ul> </li> </ul>	BET	MODERATE	International Physical	Kcal/week

	<ul style="list-style-type: none"> <li>○ One-to-one education on diet and exercise from a nurse for 5 minutes each session. Contents included consequence of obesity, dietary recommendations for weight loss, and physical activity recommendation including intensity, time, and type</li> <li>○ Fitmeter accelerometer and smartphone app to self-monitor, with or without additional financial incentives</li> </ul>		Incentive group decreased from about 0.9 to 0.5 goals accomplished per day.	Activity Questionnaire (Short Form)	
Viestar <i>et al</i> , 2018(66)	<ul style="list-style-type: none"> <li>- The VIP in Construction program includes personal health coaching, information, tools to support changes in PA and dietary behaviour</li> <li>- Two to four coaching sessions and tailored information offered during work hours including training instruction for core stability and strengthening exercises</li> <li>- “VIP in construction” toolbox (overview of company health-promoting facilities, waist-measuring tape, pedometer, BMI calculator, calorie guide, recipes, and knowledge tests)</li> </ul>	TPB HBM TTM PAPM	unknown	Short Questionnaire to Assess Health enhancing PA (SQUASH)	MVPA (min/week)
Werkman <i>et al</i> , 2010(67)	<ul style="list-style-type: none"> <li>- Five program modules over 1 year. Module 3 aimed to improve dietary and/or physical activity; participants could receive computer-tailored feedback on: physical activity, fibre, portion size. Module 1 included pedometer. Modules 4 and 5 online – participants access information on diet and physical activity behaviour.</li> </ul>	TPB TTM PAPM	MODERATE 82% used toolkit, 72% CD1, 41% CD2, 54% website and 16% interactive components	Physical Activity for the Elderly	Daily PA (min/week)
Wyke <i>et al</i> , 2019(68)	<ul style="list-style-type: none"> <li>- EuroFIT included twelve weekly 90-minute group (~15-20 men) sessions at professional football stadia. <ul style="list-style-type: none"> <li>○ Classroom-based education on topics related to increasing daily physical activity and reducing sedentary time and maintaining changes long-term, and healthy eating (including alcohol consumption)</li> <li>○ Coach-led physical activity sessions including training in aerobic, strength and flexibility exercises</li> </ul> </li> <li>- Pedometers to support self-monitoring as part of incremental pedometer-based walking program; novel pocket-worn device SitFIT) to self-monitor sedentary and non-sedentary time</li> </ul>	SCT CT	MODERATE 86% attended at least 6 of the 12 sessions, 54% attended 10 or more, 15% attended all 12 sessions	ActivPAL	Steps/day

Nil = Not identified; SCT = Social Cognitive Theory; SDT = Self Determination Theory; TTM = Transtheoretical Model; MPM = Model of Personalised Medicine; SCapT = Social Capital Theory; CT = Control theory; TPB = Theory of Planned Behaviour, FST = Family Systems Theory; SRT = Self-regulation Theory; BET = Behavioural Economic Theory; HBM = Health Belief Model; PAPM = Precaution Adoption Process Model



## Supplementary 4 (S4)

**Table S4 Study quality assessment (EPHPP)**

Primary source	Component rating						Global rating
	Selection bias	Study design	Confounders	Blinding	Data collection	Withdrawals / dropouts	Overall study quality
Aguiar <i>et al</i> , 2016(43)	Weak	Strong	Strong	Moderate	Strong	Strong	<b>Moderate</b>
Andersen <i>et al</i> , 2012(44, 45)	Weak	Strong	Strong	Moderate	Strong	Strong	<b>Moderate</b>
Ashton <i>et al</i> , 2017(46)	Weak	Strong	Strong	Moderate	Strong	Strong	<b>Moderate</b>
Galvao <i>et al</i> , 2017(47)	Moderate	Strong	Strong	Moderate	Strong	Moderate	<b>Moderate</b>
Gong <i>et al</i> , 2015(48)	Moderate	Strong	Strong	Strong	Moderate	Strong	<b>Strong</b>
Gray <i>et al</i> , 2013(49)	Weak	Strong	Strong	Weak	Strong	Strong	<b>Moderate</b>
Groeneveld <i>et al</i> , 2011(50)	Weak	Strong	Strong	Weak	Strong	Weak	<b>Moderate</b>
Hunt <i>et al</i> , 2014(51)	Weak	Strong	Strong	Weak	Strong	Weak	<b>Moderate</b>
Livingston <i>et al</i> , 2015(52, 53)	Weak	Strong	Strong	Weak	Strong	Strong	<b>Moderate</b>
Maruyama <i>et al</i> , 2010(54)	Weak	Strong	Strong	Weak	Strong	Strong	<b>Moderate</b>
McGowan <i>et al</i> , 2013(55)	Strong	Strong	Strong	Weak	Strong	Moderate	<b>Moderate</b>
Morgan <i>et al</i> , 2013(56)	Weak	Strong	Strong	Moderate	Strong	Strong	<b>Moderate</b>
Morgan <i>et al</i> , 2014(57)	Weak	Strong	Strong	Weak	Strong	Strong	<b>Moderate</b>
Morgan <i>et al</i> , 2011a(58)	Weak	Strong	Strong	Weak	Strong	Moderate	<b>Moderate</b>
Morgan <i>et al</i> , 2011b(59)	Strong	Strong	Strong	Weak	Strong	Strong	<b>Strong</b>
Morgan <i>et al</i> , 2009(60)	Weak	Strong	Strong	Strong	Strong	Strong	<b>Strong</b>
Patrick <i>et al</i> , 2011(61)	Weak	Strong	Strong	Moderate	Strong	Moderate	<b>Moderate</b>
Petrella <i>et al</i> , 2017(62)	Weak	Strong	Strong	Weak	Strong	Strong	<b>Moderate</b>
Pritchard <i>et al</i> , 1997(63)	Weak	Strong	Strong	Weak	Weak	Strong	<b>Moderate</b>
Schröder <i>et al</i> , 2018(64)	Moderate	Strong	Strong	Weak	Strong	Weak	<b>Moderate</b>
Shin <i>et al</i> , 2017(65)	Weak	Strong	Strong	Weak	Strong	Strong	<b>Moderate</b>
Viester <i>et al</i> , 2018(66)	Moderate	Strong	Strong	Moderate	Strong	Strong	<b>Strong</b>
Werkman <i>et al</i> , 2010(67)	Weak	Strong	Weak	Weak	Strong	Strong	<b>Moderate</b>
Wyke <i>et al</i> , 2019(68)	Weak	Strong	Strong	Moderate	Strong	Strong	<b>Moderate</b>