



OPEN ACCESS

COVID-19 Pandemic and Exercise (COPE) trial: a multigroup pragmatic randomised controlled trial examining effects of app-based at-home exercise programs on depressive symptoms

Eli Puterman ,¹ Benjamin Hives,¹ Nicole Mazara,¹ Nikol Grishin,¹ Joshua Webster,¹ Stacey Hutton,² Michael Stephen Koehle,^{1,3} Yan Liu,⁴ Mark R Beauchamp ¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2021-104379>).

¹School of Kinesiology, The University of British Columbia, Vancouver, British Columbia, Canada

²Canadian Sport Institute Pacific, Victoria, British Columbia, Canada

³Division of Sports Medicine, The University of British Columbia, Vancouver, British Columbia, Canada

⁴Faculty of Education, The University of British Columbia, Vancouver, British Columbia, Canada

Correspondence to

Dr Eli Puterman, The University of British Columbia, Vancouver, BC V6T 1Z4, Canada; eli.puterman@ubc.ca

Accepted 13 September 2021
Published Online First
27 September 2021



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Puterman E, Hives B, Mazara N, et al. *Br J Sports Med* 2022;**56**:546–552.

ABSTRACT

Background The number of adults across the globe with significant depressive symptoms has grown substantially during the COVID-19 pandemic. The extant literature supports exercise as a potent behaviour that can significantly reduce depressive symptoms in clinical and non-clinical populations.

Objective Using a suite of mobile applications, at-home exercise, including high intensity interval training (HIIT) and/or yoga, was completed to reduce depressive symptoms in the general population in the early months of the pandemic.

Methods A 6-week, parallel, multiarm, pragmatic randomised controlled trial was completed with four groups: (1) HIIT, (2) Yoga, (3) HIIT +yoga, and (4) waitlist control (WLC). Low active, English-speaking, non-retired Canadians aged 18–64 years were included. Depressive symptoms were measured at baseline and weekly following randomisation.

Results A total of 334 participants were randomised to one of four groups. No differences in depressive symptoms were evident at baseline. The results of latent growth modelling showed significant treatment effects in depressive symptoms for each active group compared with the WLC, with small effect sizes (ESs) in the community-based sample of participants. Treatment groups were not significantly different from each other. Effect sizes were very large (eg, week 6 ES range=–2.34 to –2.52) when restricting the analysis only to participants with high depressive symptoms at baseline.

Conclusions At-home exercise is a potent behaviour to improve mental health in adults during the pandemic, especially in those with increased levels of depressive symptoms. Promotion of at-home exercise may be a global public health target with important personal, social and economic implications as the world emerges scathed by the pandemic.

Trial registration number NCT04400279.

INTRODUCTION

Early in the COVID-19 pandemic, fears of infection, economic hardship and global stay-at-home mandates were widely predicted to worsen mental health (ie, the disruption of psychological wellbeing), including depression, anxiety and general distress.^{1–3} The prediction has been borne out, with global rates of depression and anxiety reaching 28% and 26% during the pandemic, respectively.⁴ Few

studies have investigated whether global rates of mental health concerns actually changed from pre-COVID-19 to the pandemic period. In a national poll completed 1 month into the pandemic, 10% and 23% of Canadians reported that they had high or some levels of depression, respectively, an increase from 4% and 15% reported prior to the pandemic.⁵ Twofold to threefold increases were also reported in the USA⁶ and the UK⁷ in the early months of the pandemic compared with before the pandemic.

Researchers and healthcare professionals promoted a wide range of approaches to maintain the mental health of all individuals during this pandemic, from actions individuals can take within their homes and outdoors, such as exercise, to assessment and treatment considerations that healthcare providers and institutions can implement.^{1–3} The WHO⁸ and global government agencies (eg, US Centers for Disease Control and Prevention⁹ and Public Health England)¹⁰ similarly recommended that the public engage in physical activity and exercise to attain and maintain mental health during the pandemic. These recommendations are supported by an extant literature providing compelling evidence for impactful prevention of¹¹ and reductions in¹² depressive symptoms in clinical and non-clinical populations following the adoption of physical activity programming. Many trials have also revealed the many neurobiological effects of long-term exercise in improving depressive symptoms.¹³ Yet, with the mandated closure of fitness centres and outdoor recreation sites (eg, local and state/provincial parks) at the start of the pandemic, opportunities for engaging in healthy behaviours remained limited to one's home for the most part.

Our study tested whether completing exercise at home that required little physical space or equipment would lead to reductions in depressive symptoms among Canadian adults in the spring and summer of 2020. Depressive symptoms were chosen as the trial's primary mental health outcome since loneliness/social isolation are strongly associated with depression^{14–16} and since loneliness/social isolation were particularly anticipated due to prevailing public health physical distancing mandates implemented during the early stages of the COVID-19 pandemic. Activities were completed with the use of a commercially available mobile application (app), with memberships provided free. Given that

88% of Canadians own smartphones¹⁷ and 94% of Canadians have home internet access, the use of a commercially available app that is both android and iOS compatible allowed us to test the effects of at-home exercise using app-based programs that are easily scalable to the public and widely implementable in the future, if effects are significant. The provision of free memberships and the minimal room or equipment required to complete the activities attended to some of the socioeconomic inequities that can result from at-home exercise during the pandemic, as highlighted by Sallis and colleagues.¹⁸

We partnered with a mobile application company, Down Dog (<https://www.downdogapp.com>), that has a suite of apps for a variety of activities that require little space or equipment. Two of the apps that were available for download at the start of the pandemic from Down Dog included whole body weight based high intensity interval training (HIIT) and yoga. Both HIIT¹⁹ and yoga²⁰ have been shown to be effective in improving depressive symptoms. Participants in the active groups received access to either the HIIT or yoga application or access to both applications for 6 weeks.

The primary hypothesis of our pragmatic randomised clinical trial was that completing at-home HIIT and/or yoga with the use of a free mobile application will lead to significant declines in depressive symptoms in adults over a 6-week period compared with a waitlist control (WLC). We further explored whether the benefits were unique to or stronger for HIIT, yoga or their combination by comparing each group's effects over time to one another. Finally, we tested whether the effects were more or less apparent in those with high depressive symptoms prerandomisation.

METHODS

Trial design

The COvid-19 Pandemic and Exercise (COPE) trial was a parallel, multiarm, pragmatic randomised controlled trial, with participants allocated randomly to one of four treatment groups: (1) HIIT, (2) yoga, (3) HIIT +yoga or (4) WLC. Study protocol was registered at ClinicalTrials.gov and the Open Science Framework (<https://osf.io/jbm63/>).

Participants

Low active, English-speaking, Canadians aged 18–64 years, who were not retired at study entry and had access to the internet via a mobile device or computer were eligible to participate. Activity was assessed with the validated Stanford Leisure-Time Categorical Activity Item (L-CAT).²¹ Those who scored between 1 and 3 on the L-CAT were eligible to participate since these scores represent low activity as prescribed by the American College of Sports Medicine.²² Only individuals deemed capable of performing moderate intensity physical activity were eligible to participate, as assessed with the Physical Activity Readiness Questionnaire for Everyone (PAR-Q+) and, if necessary, further assessed with the Physical Activity Readiness Medical Examination (ePARmed-X+).²³ Those hospitalised within the previous 3 months were not eligible, unless a note from their physician was provided stating their ability to participate.

Participants from across Canada were recruited through social media advertisements, including Facebook, Twitter and Instagram, reaching 21 406 unique viewers and 2731 engaged in the advertisements through shares, clicks or likes. The advertisements directed participants to the study website that provided a summary of the purpose of the study and inclusion/exclusion criteria. If interested, they were invited to email the study team

to schedule an eligibility screening phone call. A study team member completed the screening using a scripted interview with Qualtrics, where eligibility screening data were stored. On agreement, eligible and interested participants received two links by email: (1) to the Qualtrics-based consent form for electronic signature and (2) to the PAR-Q+. Interested participants who were not cleared for exercise based on the PAR-Q + or the ePARmed-X + were required to receive clearance from their family physician or our team's study physician. Following clearance, all participants completed our Qualtrics-based survey to assess the primary outcome: depressive symptoms.

Intervention

Participants randomised to one of the three treatment groups received a free 3-month membership to the mobile application version from Down Dog to access the applicable programs according to group assignment (see online supplemental materials – Methods section for details about the app). Participants in the treatment groups were asked to complete a minimum 4 weekly 20 min sessions for 6 weeks. This accumulation was based on global recommendations to complete 75 min of high/vigorous intensity activity per week (equating to 150 min of moderate intensity activity).²⁴ Seventy-five minutes of high intensity activity (for the HIIT condition) was averaged up to 80 min to translate the guidelines to 4 weekly sessions of equal length for study purposes. To ensure equivalence of weekly accumulated activity across intervention conditions, we asked participants in the yoga and HIIT +yoga conditions to also engage in four 20 min sessions per week. WLC participants were asked to remain at the same activity level as during recruitment during the 6-week study and received the free 3-month membership to the yoga and HIIT apps after the 6 weeks. To ensure anonymity on the Down Dog platform, each participant received a participant ID that was preregistered by a study team member on the Down Dog platform. This also allowed us to track their weekly progress.

Outcome

The primary outcome was depressive symptoms, measured weekly from baseline to the end of the sixth week of the trial with the 10-item Center for Epidemiological Studies – Depression Scale (CESD).²⁵ Examples of items include, 'I was bothered by things that usually don't bother me' and 'I could not get going.' Scores ranged between 0 ('Rarely or none of the time (less than 1 day)') and 3 ('Most or all of the time (5–7 days)'). Sum scores were produced (potential range from 0 to 30 (sample range: 0 to 30)). A cut-off score of 10 or above is considered significant depressive symptoms in community samples.

Sample size

Using Optimal Design Software,²⁶ 367 participants were required to detect a small effect size (ES) $\delta=0.30$ based on a two-level curvilinear growth model with power (1 - β) set at 0.80 and alpha set at 0.05 for a seven time points repeated measures design with four groups. A 25% attrition was expected over 6 weeks, thus a sample size of 490 was considered to be sufficient for the trial.

Randomisation and allocation

Sequence generation for randomisation was completed using Excel. Each member of the recruiting team received an Excel book with multiple blocks of randomised groupings. All blocks contained one of each potential grouping and a randomly

assigned number generated by the data manager (second author). The treatment groups within each block were then sorted by their randomised number, allowing for unique configurations within each block and the grouping hidden on the Excel sheet. A member of the study team would unhide the result of the randomisation and allocated the participants to their group once they had completed the baseline surveys. The data manager and principal investigator (PI) (first author) remained blind to the participants' allocations throughout the trial. The PI was blind to all randomisation until all data were prepared for analysis and initial primary analysis was completed.

Changes to the trial

On 7 August 2020, study recruitment was terminated for several reasons. First, interest in the study dropped substantially in late June, potentially due to the fact that fitness centres across Canada started to open, as did parks. Second, we wanted to keep the time frame of recruitment narrow to maintain similarity across participants in terms of impact of the pandemic. Finally, while we were very conservative with our original expected ES, a small-to-medium ES of $\delta=0.40$ or a medium ES of 0.50 could have as easily been expected based on previous meta-analyses,^{12,27} requiring 209 and 134 participants, respectively, or 278 and 179, accounting for 25% attrition. More information on these changes can be accessed here, submitted 8 August 2020: <https://osfio/a65vd/>.

Statistical approach

Means and SD, or number and percentages, were calculated for all continuous or categorical sociodemographic variables, respectively. Analyses of variance or χ^2 analyses were completed for the continuous and categorical factors, respectively, to compare group differences. Imputation, using random forest methods,²⁸ was conducted for depression symptom score when the participant did not complete all items in the survey at any week (see online supplemental statistical approach for information on imputation procedures and survey items completion rates on

online supplemental table S1). All descriptive statistics, multiple imputation and visualisations were run using R Statistical software (V.4.0.2).

All randomised participants were included in the intent-to-treat analysis using Mplus (V.7.2).²⁹ We adopted quadratic latent growth models²⁹ based on the framework of structural equation models to account for non-linear trends in CESD scores over the 6 weeks. See online supplemental materials – statistical approach section for more details of our analytic approach and the equations of the conditional quadratic latent growth models.

To examine the treatment effects in depressive symptoms on the subpopulation with high depressive symptoms initially, we restricted the sample to participants with CESD scores of ≥ 10 ²⁵ and, due to poor model fit when including the quadratic term, we used free time scores of the slope growth factor for non-linear trends (Muthén and Muthén, p. 124).²⁹ The equations are similar to those used for the quadratic latent growth model, but only I and S were estimated, and S is freely estimated with specifying the first two time points to 0 and 1.

For all analyses, we computed effect sizes at each week using Feingold's approach,³⁰ equivalent to Cohen's *d* (see online supplemental materials for additional details including *Mplus code*). Model fit indices used to ascertain model fit and fit statistics for the complete and restricted samples are included in the online supplemental materials - statistical approach section and online supplemental table S2, respectively. Maximum likelihood robust estimation was used for all the latent growth models because this type of estimator can easily handle outliers and missing data and provide more robust and accurate estimates. All codes are included in the online supplemental file 1.

RESULTS

Participants

Three hundred and ninety-six individuals were screened for the COPE trial and, based on eligibility, 334 (84%) were enrolled between 27 May 2020 and 7 August 2020 (see figure 1, consort diagram). Descriptive statistics for the sociodemographic factors




Figure 1 Consort diagram.

Table 1 Participant demographic information

Variable	All n=334	WLC n=83	HIIT n=82	Yoga n=86	HIIT+yoga n=83	Baseline comparison
Age, years (SD)	40.3 (12.4)	41.0 (12.1)	41.2 (12.7)	37.8 (12.3)	41.1 (12.6)	$F(3, 329)=1.45, p=0.23$
CESD, score (SD)	10.3 (5.78)	10.4 (5.54)	10.8 (6.23)	10.4 (5.87)	9.41 (5.43)	$F(3, 330)=0.89, p=0.45$
Women, n (%)	289 (87)	74 (89)	71 (87)	72 (84)	72 (87)	$\chi^2(3, n=334)=1.08, p=0.78$
Income, n (%)*						$\chi^2(15, n=334)=18.77, p=0.22$
0–40 000	30 (9)	4 (5)	11 (13)	9 (10)	6 (7)	
40 001–80 000	68 (20)	16 (19)	12 (15)	24 (28)	16 (19)	
80 001–120 000	66 (20)	19 (23)	14 (17)	14 (16)	19 (23)	
120 001–160 000	51 (15)	11 (13)	12 (15)	18 (21)	10 (12)	
160 000+	63 (19)	20 (24)	14 (17)	11 (13)	18 (22)	
No response	56 (17)	13 (16)	19 (23)	10 (12)	14 (17)	
Employment status†						$\chi^2(12, n=334)=6.11, p=0.91$
Full time	177 (53)	50 (60)	40 (49)	40 (47)	47 (57)	
Part time	44 (13)	10 (12)	13 (16)	13 (15)	8 (10)	
Not working	48 (14)	9 (11)	14 (17)	14 (16)	11 (13)	
Student	45 (13)	10 (12)	10 (12)	14 (16)	11 (13)	
Other	20 (6)	4 (5)	5 (6)	5 (6)	6 (7)	
Education‡						$\chi^2(12, n=334)=12.83, p=0.38$
High school or less	22 (7)	3 (4)	3 (4)	10 (12)	6 (7)	
College, trade school or certificate	52 (16)	10 (12)	12 (15)	15 (17)	15 (18)	
Bachelor or equivalent	139 (42)	37 (45)	34 (41)	36 (42)	32 (39)	
Postgraduate/professional training	118 (35)	33 (40)	31 (38)	24 (28)	30 (36)	
No response	3 (1)	0 (0)	2 (2)	1 (1)	0 (0)	
Ethnoracial identification§§						$\chi^2(6, n=334)=5.37, p=0.50$
White, n (%)	198 (59)	50 (60)	54 (66)	43 (50)	51 (61)	
Asian, n (%)	87 (26)	22 (27)	16 (20)	28 (33)	21 (25)	
Other/multiple selections/did not answer, n (%)	49 (15)	11 (13)	12 (15)	15 (17)	11 (13)	
Marital status¶						$\chi^2(9, n=334)=6.95, p=0.64$
Married, n (%)	187 (56)	49 (59)	48 (59)	43 (50)	47 (57)	
Separated/divorced/widowed, n (%)	26 (8)	7 (8)	6 (7)	5 (6)	8 (10)	
Single, n (%)	115 (34)	27 (33)	25 (30)	36 (42)	27 (33)	
Other/refused to/did not answer, n (%)	6 (2)	0 (0)	3 (4)	2 (2)	1 (1)	
LCAT						$\chi^2(6, n=334)=10.22, p=0.12$
1, n (%)	19 (6)	2 (2)	4 (5)	8 (9)	5 (6)	
2, n (%)	172 (51)	36 (43)	49 (60)	46 (53)	41 (49)	
3, n (%)	143 (43)	45 (54)	29 (35)	32 (37)	37 (45)	

Note: data are presented for all participants (all) and separated by randomisation group (waitlist control (WLC); high intensity interval training (HIIT); yoga (yoga); combination (HIIT +yoga)). All group comparisons between continuous variables were complete using one-way analyses of variance; the comparisons between categorical variables were done using χ^2 tests.

*For details of how the categories were formed, please see online supplemental table S3.

†For details of how the categories were formed, please see online supplemental table S4.

‡For details of how the categories were formed, please see online supplemental table S5.

§For details of how the categories were formed, please see online supplemental table S6.

¶For details of how the categories were formed, please see online supplemental table S7.

CESD, Centre for Epidemiological Studies – Depression Scale; LCAT, Stanford Leisure Time Activity Categorical Item.

and depressive symptoms are presented in [table 1](#). Treatment groups were not different from each other on any factor at baseline.

Adherence results

Sixty-two per cent, 64% and 75% of HIIT, yoga and HIIT +yoga participants, respectively, completed four or more sessions in the first week of the trial, with an additional 18%–29% completing between one and three workouts ([figure 2](#)). Adherence decreased during the study. While the majority of yoga and HIIT +yoga group participants were still meeting the requested four sessions per week by the end of the trial, only 40% in the HIIT group met these activity levels. Weekly completion rate for surveys by group are reported in online supplemental table S8 and weekly average minutes (and 95% CIs) of Down Dog activities completed per group are presented in online supplemental table S9. Participants in the combination group consistently engaged in more mean exercise on a weekly basis (between 71 min and 86 min

throughout the 6 weeks) than those in the yoga arm (between 69 min and 78 min) or HIIT (between 46 min and 64 min). Wait-listed participants remained underactive throughout the study, as reported on the Godin Leisure Time Exercise Questionnaire³¹ (see online supplemental materials – Methods section for more details). Online supplemental table S10 displays weekly averages and SDs of MVPA in the WLC and active groups (outside usage of the apps).

Treatment groups versus WLC

As seen in [table 2A](#) and [figure 3A](#), the WLC participants had stable depressive symptoms throughout the 6 weeks (ie, non-significant S and Q). However, HIIT and HIIT+yoga significantly reduced in depressive symptoms over time in non-linear ways ([figure 3a](#), online supplemental table S11 section A and online supplemental table S12 section A for HIIT and HIIT+yoga, respectively), whereas yoga reduced linearly over time ([figure 3a](#), online supplemental table S13 section A).




Figure 2 Exercise adherence rates by experimental condition.

Treatment effect results revealed that baseline estimates for each group were not significantly different from that of the WLC (table 2, section B1; all p 's > 0.05), whereas the growth rates over time (specifically the slopes) for HIIT and the HIIT+yoga were different from WLC (table 2, sections B2, B3). The ES for yoga (except week 1) and HIIT+yoga compared with WLC were significant with small effect sizes (range from -0.11 to -0.33), with ES getting larger over time. In the HIIT group, ES estimates were small and significant initially, though reduced in size by week 4 when they were no longer significant. See figure 3C and online supplemental table S14 for ES estimates.

Table 2 Estimates for trajectories for WLC (A) and comparisons with active treatment groups (B1–3)

	Estimate	SE	95% CI
A. Estimates, SE and 95% CI for intercept (I), slope (S) and quadratic (Q) terms for WLC			
I	10.34	0.60	9.15 to 11.49
S	-0.15	0.26	-0.65 to 0.38
Q	-0.01	0.04	-0.09 to 0.07
B. Estimates for differences between each group and WLC			
<i>B1. Differences in estimates for I</i>			
HIIT versus WLC	0.26	0.87	-1.46 to 1.97
Yoga versus WLC	-0.34	0.86	-2.02 to 1.34
HIIT+yoga versus WLC	-1.27	0.83	-2.89 to 0.35
<i>B2. Differences in estimates for S</i>			
HIIT versus WLC	-0.77	0.38	-1.51 to -0.04
Yoga versus WLC	-0.70	0.40	-1.49 to 0.08
HIIT+yoga versus WLC	-0.92	0.39	-1.69 to -0.14
<i>B3. Differences in estimates for Q</i>			
HIIT versus WLC	0.11	0.06	0.00 to 0.22
Yoga versus WLC	0.08	0.06	-0.05 to 0.20
HIIT+yoga versus WLC	0.11	0.06	-0.01 to 0.23

Results from the SEM model estimating intercept, slope and quadratic term for WLC (section A) and comparisons of these estimates with those of the three active groups (HIIT, yoga, HIIT+yoga; section B). Bold text denotes $p < 0.05$. HIIT, high intensity interval training; SEM, structural equation model; WLC, waitlist control.




Figure 3 Trajectories and effect sizes for depressive symptoms over the course of the study. Note: figure 3A shows each group's trajectories, in the full sample, while figure 3B show the trajectories for those with high (CESD score ≥ 10) levels of depressive symptoms. Figure 3C,D represent the effect sizes at each time point, for all participants and those with high levels of depressive symptoms, respectively. CESD, Center for Epidemiological Studies – Depression Scale; HIIT, high intensity interval training; WLC, waitlist control.

Comparison of treatment groups

All three groups had similar trends in decreasing depressive symptoms over the course of the study, and effect size estimates for each week were not significantly different from each other (online supplemental tables S11–13).

High depression group

When restricted to those participants with prerandomisation baseline ≥ 10 CESD, ($n=173$; mean=14.8, SD=3.98), all treatment groups had significantly greater reductions in depressive symptoms over time compared with the WLC (online supplemental table S15, section B). Rate of decrease in depressive symptoms for each group, in descending strength, was -3.39 (HIIT +yoga), -3.24 (HIIT), -3.18 (yoga) and -1.18 (WLC) (see figure 3B for trends). Within the first week, ES for each treatment group compared with WLC were significant and of small size (ES range -0.41 to -0.44) and continued to grow over the course of the study to very large when the trial was completed (ES range -2.34 to -2.51). See figure 3D and online supplemental table S13 for ES results.

DISCUSSION

Significant treatment effects in depressive symptoms were observed for participants randomised to complete HIIT or yoga, or a combination of the two, at-home using a suite of mobile applications over a 6-week period compared with WLC participants. While WLC participants' depressive symptoms remained steady throughout the 6-week period, those in the three active arms had significant declines, as hypothesised. ES at each week were small for all three active groups, with the greatest effects in the combination group. Effects were very large when the

sample was restricted to those with high depressive symptoms prior to randomisation, again with the apparent largest effects in the combined HIIT+yoga group. These differences could be attributed to the higher number of HIIT+yoga participants who completed at least four weekly workouts throughout the study, and the fact that those in the combined group completed the most minutes of activity every week. Our results highlight that providing variety to participants in pragmatic clinical trials can lead to the greatest adherence with the largest effects. This is supported by experimental research that demonstrates that those who engage in greater varieties of exercises are most likely to report more sustained physical activity.³² Our study reveals an impactful health behaviour in which adults, especially those with significant depressive symptoms, can engage that can potentially offer relief from the burden of the pandemic.

Prior to the COVID-19 pandemic, percentages of adults across the globe who were already insufficiently active were troublesome.³³ Prior to COVID-19 in Canada, only 18% of adults were physically active at recommended levels when measured with accelerometers.³⁴ Evidence suggests that these trends worsened during the early months of the pandemic, globally³⁵ and within Canada.³⁶ This is not surprising given the mandated stay-at-home orders that were implemented across all of Canada and most of the globe in the early months of the pandemic (see ref ³⁷ for a review of the timeline of restrictions in Canada during the first wave). A global call for action¹⁸ prioritised at-risk groups (eg, frontline, low-paid, healthcare system workers, laid off adults and older adults) as intervention targets for physical activity programming to reduce risk and severity of infection with COVID-19. Not included in the call for action were those at wider risk for mental health issues, and recent studies are directing attention to individuals in different countries who are at risk for pandemic-related depression. In the UK, for example, adults aged 18–34 years, women and adults with children had the greatest increases in depression during the early months of COVID-19.⁷ Similar elevated risk to women and young adults was identified in the USA, as was to those who self-identified as Hispanic, had lower education levels, were not married or had more life stressors resulting from the pandemic (eg, financial/employment loss and COVID-19 related death of a family member).⁶ In our study, the majority of participants were women (87%), nearly half were 18–39 years of age (47%) and had children at home (42%). Yet, a large majority would not be considered to have met significant economic or employment challenges, as many were employed. The extent to which exercise programming can specifically benefit adults with economic challenges remains unclear.

There are several limitations to the current study. First, while our study sought to recruit equal numbers of women and men, men did not participate at the same rate as women, likely a result of yoga appealing mostly to women.³⁸ Second, we partnered with a company with commercially available mobile apps, which allowed us to evaluate the efficacy of app-delivered programs that can be delivered at scale. While potential scalability makes our findings important, the study was not designed to control the types, difficulty or lengths of activities. Thus, it is impossible to quantify the frequency, intensity, type or time spent in specific exercises for prescriptive and clinical purposes. A final limitation is all participants were asked to complete a minimum of four sessions/week at 20 min/session, so that the total weekly HIIT would meet global recommendations for vigorous exercise.²⁴ Yet, 80 weekly minutes of moderate activity, such as yoga, do not reach the recommended 150 min of moderate intensity activity. While global recommendations are set at 75 vigorous or 150

moderate intensity minutes of physical activity per week,²⁴ there is considerable evidence suggesting that weekly physical activity even below recommended threshold levels is effective for many health issues,³⁹ including depression.⁴⁰

By 2030, the World Economic Forum projected that mental illness will account for US\$6 trillion of the annual global economic burden, accounting for more than half the burden from all non-communicable diseases.⁴¹ With the increasing prevalence of global citizens with COVID-19 related depressive symptoms, the personal, social and economic burden can be expected to be even more far-reaching and devastating. The results from the current trial suggest that health officials should continue to promote exercise to the public,¹ directing such promotion especially to those experiencing significant depressive symptoms, with the provision of low-cost or free exercise-based mobile applications to use at home as part of healthcare systems' initiatives. Free eHealth mobile applications are effective in improving medication adherence and clinic attendance for non-communicable physical diseases, leading to substantial cost-effectiveness.⁴² Perhaps, then, the free provision of exercise-based mobile applications, supported by healthcare systems, could be one method to help reduce the emerging global mental health crisis and its eventual economic burden.

What are the findings?

- ▶ This pragmatic randomised controlled trial provides evidence suggesting that at-home app-based exercise in various forms (high intensity interval training or yoga or their combination) can significantly improve depressive symptoms over a 6-week period in adults during the pandemic. Effects were strongest for those who were provided opportunities for both high intensity interval training and yoga. When the sample was restricted to only those with high baseline depressive symptoms, the weekly effects were very large.

How might it impact on clinical practice in the future?

- ▶ At-home exercise during the COVID-19 pandemic proved to be an impactful and affordable health behaviour in which adults, especially those with high depressive symptoms, can engage to bolster their mental health. In light of the long-term mental health consequences of COVID-19 with which many adults are expected to struggle, even after a return to normal, promoting and supporting programming in communities at the individual level will emerge as a necessary clinical and health policy initiative.

Correction notice This article has been corrected since it published Online First. The funding statement has been updated.

Twitter Mark R Beauchamp @BeauchampDr

Acknowledgements We would like to acknowledge Anaïs Charbonneau, Abby Cheung, Sarah de Faye, Christina Whang, Carina Wong, Tommy Yang and Rachel Zhang for their contribution to recruitment and screening of participants and to Anaïs Charbonneau, Sarah de Faye and Rachel Zhang for their contribution to data processing.

Contributors Study was conceptualised and designed by EP, MRB and NM. Implementation was completed by NM, BH, NG and JW. MSK (physician) and SH (exercise physiologist) reviewed the files of participants who did not meet clearance using our online screening tool. YL developed the analytic plan, and YL and BH completed data analyses. The first draft was completed by EP; all tables and figures were prepared by BH and reviewed by all other authors for comments.

Funding Funding for this research was provided by the Canada Research Chairs Program and Canadian Institutes of Health Research Grant # PJT 169211.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Study protocol was approved by the University of British Columbia's Behavioral Review Ethics Board (#H20-01497).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. All of the individual participant data collected during the trial after de-identification, and reported in the manuscript, will be available at <https://osf.io/g9xqp> following publication. All of the protocols, statistical analysis plan, informed consent form, analytic code, will be available 4 months following trial's end.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Eli Puterman <http://orcid.org/0000-0002-5898-2348>

Mark R Beauchamp <http://orcid.org/0000-0001-8909-3896>

REFERENCES

- Holmes EA, O'Connor RC, Perry VH, *et al.* Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry* 2020;7:547–60.
- Pfefferbaum B, North CS. Mental health and the Covid-19 pandemic. *N Engl J Med* 2020;383:510–2.
- Galea S, Merchant RM, Lurie N. The mental health consequences of COVID-19 and physical distancing: the need for prevention and early intervention. *JAMA Intern Med* 2020;180:817–8.
- Nochaiwong S, Ruengorn C, Thavorn K, *et al.* Global prevalence of mental health issues among the general population during the coronavirus disease-2019 pandemic: a systematic review and meta-analysis. *Sci Rep* 2021;11:10173.
- Mental Health Research Canada. Mental health during COVID-19 outbreak wave 1, 2020. Available: <https://static1.squarespace.com/static/5f31a311d93d0f2e28aaf04a/t/5f86800c5757437eb2b17f08/1602650133129/Full+Report+of+Findings+of+Survey+-+FINAL.pdf> [Accessed 23 Jun 2021].
- Ettman CK, Abdalla SM, Cohen GH, *et al.* Prevalence of depression symptoms in US adults before and during the COVID-19 pandemic. *JAMA Netw Open* 2020;3:e2019686.
- Pierce M, Hope H, Ford T, *et al.* Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *Lancet Psychiatry* 2020;7:883–92.
- World Health Organization. Coronavirus disease (COVID-19): staying active, 2020. Available: <https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-staying-active> [Accessed 23 Jun 2021].
- Centers for Disease Control and Prevention (CDC). COVID-19 coping with stress.
- Public Health England. Guidance for the public on the mental health and wellbeing aspects of coronavirus (COVID-19), 2020. Available: <https://www.gov.uk/government/publications/covid-19-guidance-for-the-public-on-mental-health-and-wellbeing/guidance-for-the-public-on-the-mental-health-and-wellbeing-aspects-of-coronavirus-covid-19> [Accessed 21 Jan 2021].
- Schuch FB, Vancampfort D, Firth J, *et al.* Physical activity and incident depression: a meta-analysis of prospective cohort studies. *Am J Psychiatry* 2018;175:631–48.
- Ashdown-Franks G, Firth J, Carney R, *et al.* Exercise as medicine for mental and substance use disorders: a Meta-review of the benefits for neuropsychiatric and cognitive outcomes. *Sports Med* 2020;50:151–70.
- Schuch FB, Deslandes AC, Stubbs B, *et al.* Neurobiological effects of exercise on major depressive disorder: a systematic review. *Neurosci Biobehav Rev* 2016;61:1–11.
- Beutel ME, Klein EM, Brähler E, *et al.* Loneliness in the general population: prevalence, determinants and relations to mental health. *BMC Psychiatry* 2017;17.
- Mushtaq R, Shoib S, Shah T. Psychiatric disorders and physical health? A review on the psychological aspects of loneliness. *J Clin Diagnostic Res* 2014;8:WE01–4.
- Erzen E, Çikrikci Özkan, Ö Çikrikci. The effect of loneliness on depression: a meta-analysis. *Int J Soc Psychiatry* 2018;64:427–35.
- Statistics Canada. Smartphone use and smartphone habits by gender and age group, inactive, 2021. Available: <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=2210011501> [Accessed 23 Jun 2021].
- Sallis JF, Adlaka D, Oyeyemi A, *et al.* An international physical activity and public health research agenda to inform coronavirus disease-2019 policies and practices. *J Sport Health Sci* 2020;9:328–34.
- Shepherd SO, Wilson OJ, Taylor AS, *et al.* Low-Volume high-intensity interval training in a Gym setting improves Cardio-Metabolic and psychological health. *PLoS One* 2015;10:e0139056.
- Brinsley J, Schuch F, Lederman O, *et al.* Effects of yoga on depressive symptoms in people with mental disorders: a systematic review and meta-analysis. *Br J Sports Med* 2021;55:992–1000.
- Kiernan M, Schoffman DE, Lee K, *et al.* The Stanford leisure-time activity categorical item (L-Cat): a single categorical item sensitive to physical activity changes in overweight/obese women. *Int J Obes* 2013;37:1597–602.
- American College of Sports Medicine. *Guidelines for exercise testing and prescription*. 10th Edn. Philadelphia: Wolters Kluwer, 2018. <https://www.acsm.org/read-research/books/acsm-guidelines-for-exercise-testing-and-prescription>
- Bredin SSD, Gledhill N, Jamnik VK, *et al.* PAR-Q+ and ePARmed-X+: new risk stratification and physical activity clearance strategy for physicians and patients alike. *Can Fam Physician* 2013;59:273–7.
- Bull FC, Al-Ansari SS, Biddle S, *et al.* World Health organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020;54:1451–62.
- Andresen EM, Malmgren JA, Carter WB, *et al.* Screening for depression in well older adults: evaluation of a short form of the CES-D. *Am J Prev Med* 1994;10:77–84.
- Raudenbush SW, Spybrook J, Congdon R. Optimal design plus empirical evidence (version 3.0) 2011, 2011. Available: <http://www.wtgrantfoundation.org/resources/optimal-design>
- Schuch FB, Vancampfort D, Richards J, *et al.* Exercise as a treatment for depression: a meta-analysis adjusting for publication bias. *J Psychiatr Res* 2016;77:42–51.
- Mayer M. *MissRanger: Fast Imputation of Missing Values*. R Package Version 2.1.0, 2019. Available: <https://cran.r-project.org/package=missRanger%0A>
- Muthén LK, Muthén BO. *Mplus User's Guide*. 7th edn. Los Angeles: CA, 2012.
- Feingold A. New approaches for estimation of effect sizes and their confidence intervals for treatment effects from randomized controlled trials. *Quant Method Psychol* 2019;15:96–111.
- Godin G, Shephard RJ. A simple method to assess exercise behavior in the community. *Can J Appl Sport Sci* 1985;10:141–6.
- Sylvester BD, Standage M, McEwan D, *et al.* Variety support and exercise adherence behavior: experimental and mediating effects. *J Behav Med* 2016;39:214–24.
- Guthold R, Stevens GA, Riley LM, *et al.* Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. *Lancet Glob Health* 2018;6:e1077–86.
- Statistics Canada. The daily — Canadian health measures survey: activity monitor data, 2017. Available: <https://www150.statcan.gc.ca/n1/daily-quotidien/170419/dq170419e-eng.htm> [Accessed 23 Jun 2021].
- Stockwell S, Trott M, Tully M, *et al.* Changes in physical activity and sedentary behaviours from before to during the COVID-19 pandemic lockdown: a systematic review. *BMJ Open Sport Exerc Med* 2021;7:e000960.
- Lesser IA, Nienhuis CP. The impact of COVID-19 on physical activity behavior and well-being of Canadians. *Int J Environ Res Public Health* 2020;17. doi:10.3390/ijerph17113899. [Epub ahead of print: 31 May 2020].
- Canadian Medical Association Journal News. COVID-19: A timeline of Canada's first-wave response — CMAJ News. Available: <https://cmajnews.com/2020/06/12/coronavirus-1095847/> [Accessed 23 Jun 2021].
- Park CL, Braun T, Siegel T. Who practices yoga? A systematic review of demographic, health-related, and psychosocial factors associated with yoga practice. *J Behav Med* 2015;38:460–71.
- Warburton DER, Bredin SSD. Reflections on physical activity and health: what should we recommend? *Can J Cardiol* 2016;32:495–504.
- Ravindran AV, Balneaves LG, Faulkner G, *et al.* Canadian network for mood and anxiety treatments (CANMAT) 2016 clinical guidelines for the management of adults with major depressive disorder: section 5. complementary and alternative medicine treatments. *Can J Psychiatry* 2016;61:576–87.
- Bloom DE, Cafiero ET, Jané-Llopis E. *The global economic burden of noncommunicable diseases*, 2012.
- Iribarren SJ, Cato K, Falzon L, *et al.* What is the economic evidence for mHealth? A systematic review of economic evaluations of mHealth solutions. *PLoS One* 2017;12:e0170581.

Supplemental Materials

Table of Contents

Methods.....	2
Statistical Approach.....	2
Demographic Data Coding.....	4
References.....	4
Mplus Code.....	6
All Participants Sample, Including Treatment Groups.....	6
High Depression Sample, Including Treatment Groups – Free Time Scores.....	9
Tables.....	12
Figures.....	34

Table of Tables

Table S1 - Missing Surveys at Item and Survey Levels.....	12
Table S2 - Fit Indices for the Primary SEM Models.....	13
Table S3 - Income Grouping.....	14
Table S4 - Employment Grouping.....	16
Table S5 - Education Grouping.....	19
Table S6 - Cultural Background Grouping.....	20
Table S7 - Marital Status Groupings.....	23
Table S8 - Weekly Survey Response Rate, N (%), by Experimental Grouping.....	24
Table S9 – Average Number of Minutes of DownDog Exercises Completed by Week and Experimental Group.....	25
Table S10 – Weekly Average (SD) Minutes of Moderate-to-Vigorous Physical Activity (Not Including App Usage) for Each Group.....	26
Table S11 - Estimates for Trajectories for HIIT (A) and Comparisons with WLC, Yoga and HIIT+Yoga Groups (B1-3).....	28
Table S12 - Estimates for Trajectories for HIIT+Yoga (A) and Comparisons with WLC, HIIT and Yoga Groups (B1-3).....	29
Table S13 - Estimates for Trajectories for Yoga (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1-3).....	30
Table S14 - Effect Sizes for Model with all Individuals and Model including only those with High Depressive Symptoms at Baseline.....	31
Table S15 - Estimates for Trajectories for WLC (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1 & B2) in Participants with High Levels of Depression Symptoms at Baseline.....	33

Table of Figures

Figure S1 - SEM Path Diagram for Model including all Participants.....	34
Figure S2 - SEM Path Diagram for Model including Participants with High Baseline Depressive Symptoms.....	34

Methods

Details about the Downdog app. The HIIT workout app includes >1,000 activities that require no weights or other instrumentation. At the start of each workout session, users can choose the percent of the workout that is aerobic vs. resistance training, difficulty level, type of program (e.g. Tabata, circuit training), length of session (1 min to 90min), warmup/cooldown length, interval/recovery period length, music style, and narrator's voice. The Yoga workout app includes >1,400 yoga poses, and participants can select different types of yoga practices (e.g. Ashtanga, Hatha), difficulty levels, length (5min to 90min), pace, music style, and voice. Following selection of these features in either app, a video is streamed together from the library, providing opportunities for a novel configuration each workout.

Godin Leisure Time Exercise Questionnaire. All participants completed the Godin Leisure Time Exercise Questionnaire (Courneya et al., 2004; Godin & Shephard, 1985) on a weekly basis, and indicated number and duration of light, moderate, and vigorous exercise sessions. Consistent with Courneya et al. (2004), total number of minutes of MVPA were tabulated for each participant by multiplying the number of sessions and minutes of moderate exercise and multiplying the number of sessions and minutes of vigorous exercise, and summing these two values. Consistent with recommendations by others (Tabachnick & Fidell, 2013; Wierst et al., in press), weekly out of range values (≥ 25 moderate and vigorous exercise sessions) and outliers with Z scores above 3.29 were removed.

Statistical Approach

Quadratic latent growth models. For the full sample quadratic latent growth models, we first conducted an unconditional growth model to estimate intercept (I), slope (S), and quadratic (Q)

terms. Next, we included three dummy-coded variables for the active groups, with WLC set as the comparator, to test the prespecified treatment effects of each active group on I, S, and Q.

Statistical models for quadratic latent growth model. The equations of the conditional quadratic latent growth model are provided as follows.

$$Y_{ti} = I_i + S_i * TIME_t + Q_i * TIME_t^2 + \varepsilon_{ti}$$

$$I_i = b_{00} + b_{01}G1 + b_{02}G2 + b_{03}G3 + \zeta_{0i}$$

$$S_i = b_{10} + b_{11}G1 + b_{12}G2 + b_{13}G3 + \zeta_{1i}$$

$$Q_i = b_{20} + b_{21}G1 + b_{22}G2 + b_{23}G3 + \zeta_{2i}$$

where Y_{ti} denotes the outcome for each individual participant ($i = 1, 2, \dots, n$) at each time point ($t = 0, 1, 2, \dots, 6$), $TIME_t$ denotes time scores (i.e., 0, 1, ..., 6), I_i refers to the latent intercept factor, S_i refers to the latent linear slope factor, Q_i refers to the latent quadratic slope factor, the regression coefficients for treatment conditions (G1, G2, & G3) are denoted using b , ε_{ti} denotes the individual residuals, and ζ_{0i} , ζ_{1i} , and ζ_{2i} denote the corresponding residuals for the latent growth factors (I_i , S_i , & Q_i).

Cohen's d . The general formula of the effect size measure is Cohen's $d = (b * \text{duration}) / SD$ where b denotes regression coefficient(s) of the treatment condition, duration depends on the number of weeks for a particular time point, and SD is calculated as the sum of all the corresponding variance components.

Model fit indices. Three commonly used model fit indices were used to ascertain model fit, namely a comparative fit index (CFI), the root mean square error of approximation (RMSEA), and the standardized root mean square residual (SRMR). The criteria for evaluating model fit were designated with CFI values >0.90 , and RMSEA and SRMR values $<.08$.^{32,33}

Imputation

Random forest imputation was used to impute weekly Center for Epidemiologic Studies Depression (CESD) item responses when data were missing for one or more responses to the CESD for those participants who had completed some survey data on that given week (Table S1, Non-Completely Missing Survey). Imputation was done by taking the data for each week and splitting it into those with surveys and those without. For those with surveys, any items on the CESD with missing data were imputed based on all other participant data from all weeks. Once imputation was complete, the data were rejoined with the data of those without surveys that week. This process was then repeated for each subsequent week. On occasions in which participants did not submit a completed weekly survey (Table S1, Completely Missing Surveys), no imputation was completed since the statistical approach used can handle missing data.

Demographic Data Coding

Several of the demographic variables were group together for ease of communication in tables. This was done for income (Table S3), employment (Table S4), education (Table S5), cultural background (Table S6) and marital status (Table S7).

References not Included in the Main Manuscript

- Courneya, K. S., Jones, L. W., Rhodes, R. E., & Blanchard, C. M. (2004). Effects of Different Combinations of Intensity Categories on Self-Reported Exercise. *Research Quarterly for Exercise and Sport*, 75(4), 429–433. <https://doi.org/10.1080/02701367.2004.10609176>
- Godin, G., & Shephard, R. J. (1985). A simple method to assess exercise behavior in the community. *Canadian Journal of Applied Sport Sciences. Journal Canadien Des Sciences Appliquees Au Sport*, 10(3), 141–146.
- Tabachnick, B. G., & Fidell, L. S. (2013). *Using Multivariate Statistics* (6th editio). Pearson.
- Wierst, C., Zumbo, B., Rhodes, R. E., Faulkner, G. E., & Beauchamp, M. R. (in press). An

examination of Dweck's psychological needs model in relation to exercise-related well-being. *Journal of Sport and Exercise Psychology*, 1–12. <https://doi.org/10.1123/jsep.2021-0026>

Mplus Code***All Participants Sample, Including Treatment Groups***

VARIABLE: NAMES ARE id cond g1 g2 g3 male age

CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6 CESD12;

MISSING=ALL(999);

USEVARIABLES ARE g1 g2 g3

CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6;

ANALYSIS: ESTIMATOR = MLR;

STITERATIONS=20000;

ITERATION = 40000;

MODEL:

i s q | CESD0@0 CESD1@1 CESD2@2 CESD3@3 CESD4@4 CESD5@5 CESD6@6;

i s q ON g1 g2 g3;

s ON g1 (bs1);

s ON g2 (bs2);

s ON g3 (bs3);

q ON g1 (bq1);

q ON g2 (bq2);

q ON g3 (bq3);

i(v0);

CESD0 - CESD6 (r0-r6);

MODEL CONSTRAINT:

NEW(g1dt1 g1dt2 g1dt3 g1dt4 g1dt5 g1dt6);

$$g1dt1 = (bs1*1 + bq1*1)/\sqrt{(v0+r0/2+r1/2)};$$

$$g1dt2 = (bs1*2 + bq1*4)/\sqrt{(v0+r0/3+r1/3+r2/3)};$$

$$g1dt3 = (bs1*3 + bq1*9)/\sqrt{(v0+r0/4+r1/4+r2/4+r3/4)};$$

$$g1dt4 = (bs1*4 + bq1*16)/\sqrt{(v0+r0/5+r1/5+r2/5+r3/5+r4/5)};$$

$$g1dt5 = (bs1*5 + bq1*25)/\sqrt{(v0+r0/6+r1/6+r2/6+r3/5+r4/6+r5/6)};$$

$$g1dt6 = (bs1*6 + bq1*36)/\sqrt{(v0+r0/7+r1/7+r2/7+r3/7+r4/7+r5/7+r6/7)};$$

NEW(g2dt1 g2dt2 g2dt3 g2dt4 g2dt5 g2dt6);

$$g2dt1 = (bs2*1 + bq2*1)/\sqrt{(v0+r0/2+r1/2)};$$

$$g2dt2 = (bs2*2 + bq2*4)/\sqrt{(v0+r0/3+r1/3+r2/3)};$$

$$g2dt3 = (bs2*3 + bq2*9)/\sqrt{(v0+r0/4+r1/4+r2/4+r3/4)};$$

$$g2dt4 = (bs2*4 + bq2*16)/\sqrt{(v0+r0/5+r1/5+r2/5+r3/5+r4/5)};$$

$$g2dt5 = (bs2*5 + bq2*25)/\sqrt{(v0+r0/6+r1/6+r2/6+r3/5+r4/6+r5/6)};$$

$$g2dt6 = (bs2*6 + bq2*36)/\sqrt{(v0+r0/7+r1/7+r2/7+r3/7+r4/7+r5/7+r6/7)};$$

NEW(g3dt1 g3dt2 g3dt3 g3dt4 g3dt5 g3dt6);

$$g3dt1 = (bs3*1 + bq3*1)/\sqrt{(v0+r0/2+r1/2)};$$

$$g3dt2 = (bs3*2 + bq3*4)/\sqrt{(v0+r0/3+r1/3+r2/3)};$$

$$g3dt3 = (bs3*3 + bq3*9)/\sqrt{(v0+r0/4+r1/4+r2/4+r3/4)};$$

$$g3dt4 = (bs3*4 + bq3*16)/\sqrt{(v0+r0/5+r1/5+r2/5+r3/5+r4/5)};$$

$$g3dt5 = (bs3*5 + bq3*25)/\sqrt{(v0+r0/6+r1/6+r2/6+r3/5+r4/6+r5/6)};$$

$$g3dt6 = (bs3*6 + bq3*36)/\sqrt{(v0+r0/7+r1/7+r2/7+r3/7+r4/7+r5/7+r6/7)};$$

OUTPUT: SAMPSTAT CINTERVAL STANDARDIZED RESIDUAL MODINDICES (3.84);

High Depression Sample, Including Treatment Groups – Free Time Scores

VARIABLE: NAMES ARE id cond g1 g2 g3 male age

CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6 CESD12;

MISSING=ALL(999);

USEVARIABLES ARE g1 g2 g3 CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6;

ANALYSIS: ESTIMATOR = MLR;

STITERATIONS=5000;

ITERATION = 20000;

MODEL:

i s | CESD0@0 CESD1@1 CESD2* CESD3* CESD4* CESD5* CESD6*;

i s ON g1 g2 g3;

s ON g1 (bs1);

s ON g2 (bs2);

s ON g3 (bs3);

i(v0);

CESD1 - CESD6 (r1-r6);

CESD0@0;

CESD1 WITH CESD2-CESD4;

CESD2 WITH CESD3-CESD5;

CESD3 WITH CESD4-CESD6;

CESD4 WITH CESD5-CESD6;

CESD5 WITH CESD6;

MODEL CONSTRAINT:

NEW(g1dt1 g1dt2 g1dt3 g1dt4 g1dt5 g1dt6);

$g1dt1 = (bs1 * 1) / \sqrt{v0 + r1}$;

$g1dt2 = (bs1 * 2) / \sqrt{v0 + r1/2 + r2/2}$;

$g1dt3 = (bs1 * 3) / \sqrt{v0 + r1/3 + r2/3 + r3/3}$;

$g1dt4 = (bs1 * 4) / \sqrt{v0 + r1/4 + r2/4 + r3/4 + r4/4}$;

$g1dt5 = (bs1 * 5) / \sqrt{v0 + r1/5 + r2/5 + r3/5 + r4/5 + r5/5}$;

$g1dt6 = (bs1 * 6) / \sqrt{v0 + r1/6 + r2/6 + r3/6 + r4/6 + r5/6 + r6/6}$;

NEW(g2dt1 g2dt2 g2dt3 g2dt4 g2dt5 g2dt6);

$g2dt1 = (bs2 * 1) / \sqrt{v0 + r1}$;

$g2dt2 = (bs2 * 2) / \sqrt{v0 + r1/2 + r2/2}$;

$g2dt3 = (bs2 * 3) / \sqrt{v0 + r1/3 + r2/3 + r3/3}$;

$g2dt4 = (bs2 * 4) / \sqrt{v0 + r1/4 + r2/4 + r3/4 + r4/4}$;

$g2dt5 = (bs2 * 5) / \sqrt{v0 + r1/5 + r2/5 + r3/5 + r4/5 + r5/5}$;

$g2dt6 = (bs2 * 6) / \sqrt{v0 + r1/6 + r2/6 + r3/6 + r4/6 + r5/6 + r6/6}$;

NEW(g3dt1 g3dt2 g3dt3 g3dt4 g3dt5 g3dt6);

$g3dt1 = (bs3 * 1) / \sqrt{v0 + r1}$;

$$g3dt2 = (bs3*2)/\sqrt{(v0+r1/2+r2/2)};$$

$$g3dt3 = (bs3*3)/\sqrt{(v0+r1/3+r2/3+r3/3)};$$

$$g3dt4 = (bs3*4)/\sqrt{(v0+r1/4+r2/4+r3/4+r4/4)};$$

$$g3dt5 = (bs3*5)/\sqrt{(v0+r1/5+r2/5+r3/5+r4/5+r5/5)};$$

$$g3dt6 = (bs3*6)/\sqrt{(v0+r1/6+r2/6+r3/6+r4/6+r5/6+r6/6)};$$

OUTPUT: SAMPSTAT CINTERVAL STANDARDIZED RESIDUAL;

Tables*Table S1 - Missing Surveys at Item and Survey Levels.*

Week	Non-missing Survey (All items complete)	Non-Completely Missing Survey (Missing ≥ 1 items)	Completely Missing Surveys
0	327	7	0
1	306	9	19
2	299	5	30
3	290	2	42
4	287	6	41
5	276	6	52
6	284	6	44

Table S2 - Fit Indices for the Primary SEM Models.

Measure	All Participants - Quadratic Time	High Depression – Free Time Scores
Root Mean Square Error of Approximation, Est. (.90CI)	0.049 (0.027, 0.069)	0.042 (0.000, 0.081)
CFI	0.983	0.991
Standardized Root Mean Square Residual	0.026	0.044

Table S3 - Income Grouping.

Income Values	Recoded
1 - 10,000 10,001 - 20,000 20,001 - 30,000 30,001 - 40,000	0 – 40,000
40,001 - 50,000, 50,001 - 60,000, 60,001 - 70,000, 70,001 - 80,000	40,001 – 80,000
80,001 - 90,000, 90,001 - 100,000, 100,001 - 110,000, 110,001 - 120,000	80,001 – 120,000
120,001 - 130,000, 130,001 - 140,000, 140,001 - 150,000, 150,001 - 160,000	120,001 – 160,000
160,001 - 170,000, 170,001 - 180,000, 180,001 - 190,000, 190,001 - 200,000, 200,001 - 210,000, 210,001 - 220,000,	160,000 +

220,001 - 230,000, 230,001 - 240,000, 240,001 - 250,000, 250,001+	
Do not know / prefer not to answer, [Missing]	No response

Table S4 - Employment Grouping.

Current Employment	Other (Text Response)	Coded Value
Other	Full time work with one furlough day due to covid	Full Time
Other	teacher in spring semester	Full Time
Other	Teacher summer vacation	Full Time
Working full-time		Full Time
Working full-time,Other	on summer vacation	Full Time
Working full-time,Self-employed		Full Time
Working full-time,Self-employed ,Other	I'm working full time at a company, but also self-employed teaching (Not full-time-self-employed)	Full Time
Homemaker		Not Working
Looking for work; unemployed		Not Working
Looking for work; unemployed,Homemaker		Not Working
Looking for work; unemployed,Other	housewife	Not Working
Looking for work; unemployed,Other	working - unpaid	Not Working
Looking for work; unemployed,Temporarily laid off		Not Working
Maternity or sick leave (volunteered)		Not Working
Other	Leave of absence	Not Working

Other	Self employed but no work available	Not Working
Other	Stay at home parent	Not Working
Self-employed ,Looking for work; unemployed,Permanently disabled (volunteered),Other	Home based business not in market due to covid	Not Working
Self-employed ,Temporarily laid off		Not Working
Temporarily laid off		Not Working
Temporarily laid off,Maternity or sick leave (volunteered)		Not Working
[Missing]		Other
Don't know/not sure		Other
Other	Will be laid off during this study	Other
Other		Other
Prefer not to say		Other
Self-employed		Other
Working part-time		Part Time
Working part-time,Homemaker		Part Time
Working part-time,Other	on CEWS	Part Time
Working part-time,Self-employed		Part Time
Working part-time,Self-employed ,Homemaker		Part Time
Full-time student		Student

Looking for work; unemployed,Full-time student		Student
Looking for work; unemployed,Part-time student		Student
Looking for work; unemployed,Temporarily laid off,Part-time student		Student
Part-time student		Student
Self-employed ,Full-time student		Student
Temporarily laid off,Full-time student		Student
Temporarily laid off,Part-time student		Student
Working full-time,Full-time student		Student
Working full-time,Part-time student		Student
Working part-time,Full-time student		Student
Working part-time,Full-time student,Other	Support from parents	Student
Working part-time,Part-time student		Student

Table S5 - Education Grouping.

Educational Status	Recoded
High school diploma or a high school equivalency certificate	High School or less
College, CEGEP or other non-university certificate or diploma (other than trades certificates or diplomas),	College, trade school or certificate
Trade certificate or diploma,	College, trade school or certificate
University certificate or diploma below the bachelor's level	College, trade school or certificate
Bachelor's degree (e.g. B.A., B.Sc., LL.B.)	Bachelor or equivalent
University certificate, diploma, degree above the bachelor's level	Postgraduate / professional training
Prefer not to answer,	No response
[Missing]	No response

Note: "Less than high school diploma or its equivalent" was an option, but no participant selected it.

Table S6 - Cultural Background Grouping.

Cultural Background	Other (Text Response)	Coded Value
Chinese		Asian
Chinese,Filipino		Asian
Chinese,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Asian
Filipino		Asian
Filipino,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Asian
Japanese		Asian
Korean		Asian
Other - please specify	Taiwanese	Asian
South Asian (e.g., East Indian, Pakistani, Sri Lankan, etc)		Asian
South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Asian
Aboriginal decent (e.g., North American Indian, Métis or Inuit (Eskimo))		Other
Aboriginal decent (e.g., North American Indian, Métis or Inuit (Eskimo)),Black (e.g., African, Haitian, Jamaican, Somali, etc)		Other
Aboriginal decent (e.g., North American Indian, Métis or Inuit (Eskimo)),White		Other

Arab		Other
Black (e.g., African, Haitian, Jamaican, Somali, etc)		Other
Latin American		Other
Other - please specify		Other
Other - please specify	european / mixed	Other
Other - please specify	Doesn't let you choose more than one. Aboriginal and European.	Other
Other - please specify	Afghan	Other
Other - please specify	Austrian	Other
South Asian (e.g., East Indian, Pakistani, Sri Lankan, etc),Latin American		Other
West Asian (e.g., Iranian, Afghan, etc)		Other
White,Arab		Other
White,Chinese		Other
White,Chinese,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Other
White,Filipino		Other
White,Japanese		Other
White,Latin American		Other
White,Other - please specify	Jewish	Other

White,Other - please specify	European	Other
White,Other - please specify	Ashkenazi Jewish	Other
White,Other - please specify	Scandinavian	Other
White,Other - please specify	Dutch	Other
White,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Other
Other - please specify	Italian	White
Other - please specify	IRISH	White
White		White

Note: As cultural backgrounds were asked as check boxes, many participant selected multiple options, making it difficult to define groups. Therefore, participants were split into White, Asian and Other, as White and Asian participants made up a large majority of the sample.

Table S7 - Marital Status Groupings.

Marital Status	Recoded
Married	Married
Living common-law	Married
Separated	No Longer Married
Divorced	No Longer Married
Widowed	No Longer Married
Single, never married	Single
Prefer not to answer	Other
[Missing]	Other

Table S8 - Weekly Survey Response Rate, N (%), by Experimental Grouping.

	Week						
	0	1	2	3	4	5	6
WLC	83 (100%)	80 (96%)	77 (93%)	80 (96%)	76 (92%)	77 (93%)	74 (89%)
HIIT	82 (100%)	77 (94%)	76 (93%)	69 (84%)	72 (88%)	64 (78%)	68 (83%)
Yoga	86 (100%)	83 (97%)	76 (88%)	74 (86%)	76 (88%)	75 (87%)	67(78%)
HIIT+Yoga	83 (100%)	75 (90%)	75 (90%)	69 (83%)	69 (83%)	66 (80%)	84 (82%)
	Week						
	0	1	2	3	4	5	6
WLC	83 (100%)	80 (96%)	77 (93%)	80 (96%)	76 (92%)	77 (93%)	74 (89%)
HIIT	82 (100%)	77 (94%)	76 (93%)	69 (84%)	72 (88%)	64 (78%)	68 (83%)
Yoga	86 (100%)	83 (97%)	76 (88%)	74 (86%)	76 (88%)	75 (87%)	67(78%)
HIIT+Yoga	83 (100%)	75 (90%)	75 (90%)	69 (83%)	69 (83%)	66 (80%)	84 (82%)

Table S9 – Average Number of Minutes of DownDog Exercises Completed by Week and Experimental Group

Week	HIIT	Yoga	Combo
1	63.95 (29.73)	73.13 (43.43)	80.72 (48.48)
2	64.44 (35.03)	78.43 (46.70)	86.04 (53.80)
3	59.39 (35.49)	76.76 (53.98)	78.31 (45.19)
4	58.90 (36.08)	73.44 (59.71)	74.20 (55.49)
5	53.15 (36.85)	68.57 (50.81)	72.86 (56.29)
6	46.11 (39.02)	68.60 (57.62)	70.97 (62.29)

Table S10 – Weekly Average (SD) Minutes of Moderate-to-Vigorous Physical Activity (Not Including App Usage) for Each Group

Group	Week	Moderate	Vigorous	MVPA
Control	0	16.73 (33.25)	0.00 (0.00)	18.51 (34.53)
Control	1	40.71 (81)	7.49 (21.95)	49.04 (90.77)
Control	2	42.68 (63.32)	9.32 (29.21)	52.57 (72.50)
Control	3	44.32 (60.7)	4.34 (15.37)	47.15 (65.95)
Control	4	35.42 (48.19)	7.47 (21.26)	43.38 (57.89)
Control	5	44.17 (67.25)	6.1 (18.66)	49.91 (72.59)
Control	6	71.67 (83.48)	6.16 (16.89)	74.52 (81.26)
HIIT	0	19.72 (34.26)	0.30 (2.46)	19.06 (34.79)
HIIT	1	40.00 (52.03)	6.38 (19.11)	48.49 (60.39)
HIIT	2	45.36 (59.22)	15.40 (35.00)	61.88 (74.94)
HIIT	3	45.17 (66.58)	12.71 (26.82)	59.12 (79.29)
HIIT	4	59.25 (82.2)	14.32 (31.79)	71.29 (89.25)
HIIT	5	41.86 (55.51)	15.26 (31.00)	57.02 (73.43)
HIIT	6	49.33 (80.68)	13.71 (29.41)	59.69 (88.17)
Yoga	0	26.86 (38.26)	1.01 (4.25)	24.57 (36.49)
Yoga	1	56.88 (89.41)	11.46 (31.36)	68.66 (107.56)
Yoga	2	44.42 (58.8)	7.17 (21.77)	52.58 (67.24)
Yoga	3	47.84 (60.56)	14.18 (25.82)	58.15 (69.24)
Yoga	4	56.94 (68.03)	16.54 (33.89)	73.38 (81.59)
Yoga	5	73.46 (106.01)	11.42 (24.41)	85.07 (116.04)

Yoga	6	51.58 (72.44)	15.91 (30.62)	66.75 (92.63)
Combo	0	18.87 (35.98)	0.99 (4.11)	20.46 (37.72)
Combo	1	46.15 (97.01)	10.73 (29.82)	58.07 (103.51)
Combo	2	32.46 (45.73)	10.76 (27.46)	43.43 (54.73)
Combo	3	56.52 (70.15)	13.52 (25.7)	67.59 (74.84)
Combo	4	49.41 (68.91)	17.70 (33.67)	65.78 (85.26)
Combo	5	66.25 (99.03)	10.79 (26.15)	75.20 (112.14)
Combo	6	53.49 (74.67)	12.66 (29.28)	59.75 (78.46)

Table S11 - Estimates for Trajectories for HIIT (A) and Comparisons with WLC, Yoga and HIIT+Yoga Groups (B1-3).

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I; estimated baseline), slope (S; time), and quadratic (Q; time²) terms for HIIT</u>			
I	10.60	0.64	9.34, 11.86
S	-0.93	0.27	-1.45, -0.40
Q	0.10	0.04	0.03, 0.18
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
WLC vs HIIT	-0.26	0.87	-1.97, 1.46
Yoga vs HIIT	-0.59	0.89	-2.34, 1.16
HIIT+Yoga vs HIIT	-1.53	0.87	-3.23, 0.17
<u>B2. Differences in estimates for S</u>			
WLC vs HIIT	0.77	0.38	0.04, 1.51
Yoga vs HIIT	0.07	0.40	-0.72, 0.86
HIIT+Yoga vs HIIT	-0.14	0.40	-0.93, 0.65
<u>B3. Differences in estimates for Q</u>			
WLC vs HIIT	-0.11	0.06	-0.22, 0.00
Yoga vs HIIT	-0.03	0.06	-0.15, 0.08
HIIT+Yoga vs HIIT	0.01	0.06	-0.11, 0.12

Note: Results from the SEM. model estimating intercept, slope, and quadratic term for HIIT group (Section A) and comparisons of these estimates with those of the three other groups (WLC, Yoga, HIIT+Yoga; Section B). Bold text denotes $p < .05$

Table S12 - Estimates for Trajectories for HIIT+Yoga (A) and Comparisons with WLC, HIIT and Yoga Groups (B1-3).

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I; estimated baseline), slope (S; time), and quadratic (Q; time²) terms for HIIT+Yoga</u>			
I	9.07	0.58	7.93, 10.20
S	-1.07	0.30	-1.65, -0.48
Q	0.11	0.05	0.02, 0.19
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
WLC vs HIIT+Yoga	1.27	0.83	-0.35, 2.89
HIIT vs HIIT+Yoga	1.53	0.87	-0.17, 3.23
Yoga vs HIIT+Yoga	0.94	0.85	-0.72, 2.59
<u>B2. Differences in estimates for S</u>			
WLC vs HIIT+Yoga	0.92	0.39	0.14, 1.69
HIIT vs HIIT+Yoga	0.14	0.40	-0.65, 0.93
Yoga vs HIIT+Yoga	0.21	0.42	-0.62, 1.04
<u>B3. Differences in estimates for Q</u>			
WLC vs HIIT+Yoga	-0.11	0.06	-0.23, 0.01
HIIT vs HIIT+Yoga	-0.01	0.06	-0.12, 0.11
Yoga vs HIIT+Yoga	-0.04	0.06	-0.16, 0.09

Note: Results from the SEM model estimating intercept, slope, and quadratic term for HIIT+Yoga group (Section A) and comparisons of these estimates with those of the three other groups (WLC, HIIT, Yoga; Section B). Bold text denotes $p < .05$

Table S13 - Estimates for Trajectories for Yoga (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1-3).

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I; estimated baseline), slope (S; time), and quadratic (Q; time²) terms for Yoga</u>			
I	10.01	0.62	8.78, 11.23
S	-0.86	0.31	-1.45, -0.26
Q	0.07	0.05	-0.02, 0.16
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
WLC vs Yoga	0.34	0.86	-1.34, 2.02
HIIT vs Yoga	0.59	0.89	-1.16, 2.34
HIIT+Yoga vs Yoga	-0.94	0.85	-2.59, 0.72
<u>B2. Differences in estimates for S</u>			
WLC vs Yoga	0.70	0.40	-0.08, 1.49
HIIT vs Yoga	-0.07	0.40	-0.86, 0.72
HIIT+Yoga vs Yoga	-0.21	0.422	-1.04, 0.62
<u>B3. Differences in estimates for Q</u>			
WLC vs Yoga	-0.08	0.06	-0.20, 0.05
HIIT vs Yoga	0.03	0.06	-0.08, 0.15
HIIT+Yoga vs Yoga	0.04	0.06	-0.09, 0.16

Note: Results from the SEM. model estimating intercept, slope, and quadratic term for HIIT group (Section A) and comparisons of these estimates with those of the three other groups (WLC, HIIT, HIIT+Yoga; Section B). Bold text denotes $p < .05$

Table S14 - Effect Sizes for Model with all Individuals and Model including only those with High Depressive Symptoms at Baseline.

All Participants						
Time	HIIT		Yoga		HIIT+Yoga	
	Effect Size	95% CI	Effect Size	95% CI	Effect Size	95% CI
Week 1	-0.12	[-0.23, -0.01]	-0.11	[-0.23, 0.01]	-0.14	[-0.26, -0.03]
Week 2	-0.20	[-0.39, -0.01]	-0.20	[-0.39, 0.00]	-0.24	[-0.44, -0.05]
Week 3	-0.24	[-0.48, 0.00]	-0.25	[-0.49, -0.02]	-0.31	[-0.54, -0.07]
Week 4	-0.24	[-0.50, 0.02]	-0.28	[-0.53, -0.04]	-0.33	[-0.58, -0.08]
Week 5	-0.21	[-0.48, 0.06]	-0.29	[-0.52, -0.06]	-0.31	[-0.55, -0.07]
Week 6	-0.14	[-0.44, 0.16]	-0.27	[-0.51, -0.04]	-0.25	[-0.51, 0.00]
Subpopulation with High Depressive Symptoms						
Time	HIIT		Yoga		HIIT+Yoga	
	Effect Size	95% CI	Effect Size	95% CI	Effect Size	95% CI
Week 1	-0.41	[-0.69, -0.14]	-0.40	[-0.67, -0.14]	-0.44	[-0.71, -0.18]
Week 2	-0.83	[-1.38, -0.28]	-0.80	[-1.33, -0.28]	-0.89	[-1.43, -0.35]

Week 3	-1.20	[-2.01, -0.40]	-1.17	[-1.94, -0.39]	-1.29	[-2.07, -0.51]
Week 4	-1.57	[-2.62, -0.52]	-1.52	[-2.53, -0.51]	-1.68	[-2.70, -0.67]
Week 5	-1.94	[-3.21, -0.66]	-1.88	[-3.11, -0.65]	-2.08	[-3.32, -0.84]
Week 6	-2.34	[-3.87, -0.81]	-2.27	[-3.74, -0.80]	-2.51	[-4.00, -1.01]

Note: All effect sizes are compared to the WLC group.

Table S15 - Estimates for Trajectories for WLC (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1 & B2) in Participants with High Levels of Depression Symptoms at Baseline.

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I) and slope (S) terms for WLC</u>			
I	14.41	0.56	13.32, 15.51
S	-1.18	0.44	-2.05, -0.31
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
HIIT vs WLC	1.04	0.85	-0.62, 2.70
Yoga vs WLC	0.52	0.81	-1.06, 2.10
HIIT+Yoga vs WLC	-0.26	0.83	-1.89, 1.38
<u>B2. Differences in estimates for S</u>			
HIIT vs WLC	-2.06	0.68	-3.39, -0.73
Yoga vs WLC	-2.00	0.65	-3.28, -0.72
HIIT+Yoga vs WLC	-2.21	0.67	-3.52, -0.90

Note: Results from the SEM model, including only those with high levels of depressive symptoms at baseline, estimating intercept and slope for waitlist control (Section A) and comparisons of these estimates with those of the three active groups (HIIT, Yoga, HIIT+Yoga; Section B). Bold text denotes $p < .05$

Figures

Figure S1 - SEM Path Diagram for Model including all Participants.





Figure S2 - SEM Path Diagram for Model including Participants with High Baseline Depressive Symptoms.



Supplemental Materials

Table of Contents

Methods.....	2
Statistical Approach.....	2
Demographic Data Coding.....	4
References.....	4
Mplus Code.....	6
All Participants Sample, Including Treatment Groups.....	6
High Depression Sample, Including Treatment Groups – Free Time Scores.....	9
Tables.....	12
Figures.....	34

Table of Tables

Table S1 - Missing Surveys at Item and Survey Levels.....	12
Table S2 - Fit Indices for the Primary SEM Models.....	13
Table S3 - Income Grouping.....	14
Table S4 - Employment Grouping.....	16
Table S5 - Education Grouping.....	19
Table S6 - Cultural Background Grouping.....	20
Table S7 - Marital Status Groupings.....	23
Table S8 - Weekly Survey Response Rate, N (%), by Experimental Grouping.....	24
Table S9 – Average Number of Minutes of DownDog Exercises Completed by Week and Experimental Group.....	25
Table S10 – Weekly Average (SD) Minutes of Moderate-to-Vigorous Physical Activity (Not Including App Usage) for Each Group.....	26
Table S11 - Estimates for Trajectories for HIIT (A) and Comparisons with WLC, Yoga and HIIT+Yoga Groups (B1-3).....	28
Table S12 - Estimates for Trajectories for HIIT+Yoga (A) and Comparisons with WLC, HIIT and Yoga Groups (B1-3).....	29
Table S13 - Estimates for Trajectories for Yoga (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1-3).....	30
Table S14 - Effect Sizes for Model with all Individuals and Model including only those with High Depressive Symptoms at Baseline.....	31
Table S15 - Estimates for Trajectories for WLC (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1 & B2) in Participants with High Levels of Depression Symptoms at Baseline.....	33

Table of Figures

Figure S1 - SEM Path Diagram for Model including all Participants.....	34
Figure S2 - SEM Path Diagram for Model including Participants with High Baseline Depressive Symptoms.....	34

Methods

Details about the Downdog app. The HIIT workout app includes >1,000 activities that require no weights or other instrumentation. At the start of each workout session, users can choose the percent of the workout that is aerobic vs. resistance training, difficulty level, type of program (e.g. Tabata, circuit training), length of session (1 min to 90min), warmup/cooldown length, interval/recovery period length, music style, and narrator's voice. The Yoga workout app includes >1,400 yoga poses, and participants can select different types of yoga practices (e.g. Ashtanga, Hatha), difficulty levels, length (5min to 90min), pace, music style, and voice. Following selection of these features in either app, a video is streamed together from the library, providing opportunities for a novel configuration each workout.

Godin Leisure Time Exercise Questionnaire. All participants completed the Godin Leisure Time Exercise Questionnaire (Courneya et al., 2004; Godin & Shephard, 1985) on a weekly basis, and indicated number and duration of light, moderate, and vigorous exercise sessions. Consistent with Courneya et al. (2004), total number of minutes of MVPA were tabulated for each participant by multiplying the number of sessions and minutes of moderate exercise and multiplying the number of sessions and minutes of vigorous exercise, and summing these two values. Consistent with recommendations by others (Tabachnick & Fidell, 2013; Wierst et al., in press), weekly out of range values (≥ 25 moderate and vigorous exercise sessions) and outliers with Z scores above 3.29 were removed.

Statistical Approach

Quadratic latent growth models. For the full sample quadratic latent growth models, we first conducted an unconditional growth model to estimate intercept (I), slope (S), and quadratic (Q)

terms. Next, we included three dummy-coded variables for the active groups, with WLC set as the comparator, to test the prespecified treatment effects of each active group on I, S, and Q.

Statistical models for quadratic latent growth model. The equations of the conditional quadratic latent growth model are provided as follows.

$$Y_{ti} = I_i + S_i * TIME_t + Q_i * TIME_t^2 + \varepsilon_{ti}$$

$$I_i = b_{00} + b_{01}G1 + b_{02}G2 + b_{03}G3 + \zeta_{0i}$$

$$S_i = b_{10} + b_{11}G1 + b_{12}G2 + b_{13}G3 + \zeta_{1i}$$

$$Q_i = b_{20} + b_{21}G1 + b_{22}G2 + b_{23}G3 + \zeta_{1i}$$

where Y_{ti} denotes the outcome for each individual participant ($i = 1, 2, \dots, n$) at each time point ($t = 0, 1, 2, \dots, 6$), $TIME_t$ denotes time scores (i.e., 0, 1, ..., 6), I_i refers to the latent intercept factor, S_i refers to the latent linear slope factor, Q_i refers to the latent quadratic slope factor, the regression coefficients for treatment conditions (G1, G2, & G3) are denoted using b , ε_{ti} denotes the individual residuals, and ζ_{0i} , ζ_{1i} , and ζ_{1i} denote the corresponding residuals for the latent growth factors (I_i , S_i , & Q_i).

Cohen's d . The general formula of the effect size measure is Cohen's $d = (b * \text{duration}) / SD$ where b denotes regression coefficient(s) of the treatment condition, duration depends on the number of weeks for a particular time point, and SD is calculated as the sum of all the corresponding variance components.

Model fit indices. Three commonly used model fit indices were used to ascertain model fit, namely a comparative fit index (CFI), the root mean square error of approximation (RMSEA), and the standardized root mean square residual (SRMR). The criteria for evaluating model fit were designated with CFI values >0.90 , and RMSEA and SRMR values $<.08$.^{32,33}

Imputation

Random forest imputation was used to impute weekly Center for Epidemiologic Studies Depression (CESD) item responses when data were missing for one or more responses to the CESD for those participants who had completed some survey data on that given week (Table S1, Non-Completely Missing Survey). Imputation was done by taking the data for each week and splitting it into those with surveys and those without. For those with surveys, any items on the CESD with missing data were imputed based on all other participant data from all weeks. Once imputation was complete, the data were rejoined with the data of those without surveys that week. This process was then repeated for each subsequent week. On occasions in which participants did not submit a completed weekly survey (Table S1, Completely Missing Surveys), no imputation was completed since the statistical approach used can handle missing data.

Demographic Data Coding

Several of the demographic variables were group together for ease of communication in tables. This was done for income (Table S3), employment (Table S4), education (Table S5), cultural background (Table S6) and marital status (Table S7).

References not Included in the Main Manuscript

- Courneya, K. S., Jones, L. W., Rhodes, R. E., & Blanchard, C. M. (2004). Effects of Different Combinations of Intensity Categories on Self-Reported Exercise. *Research Quarterly for Exercise and Sport*, 75(4), 429–433. <https://doi.org/10.1080/02701367.2004.10609176>
- Godin, G., & Shephard, R. J. (1985). A simple method to assess exercise behavior in the community. *Canadian Journal of Applied Sport Sciences. Journal Canadien Des Sciences Appliquees Au Sport*, 10(3), 141–146.
- Tabachnick, B. G., & Fidell, L. S. (2013). *Using Multivariate Statistics* (6th editio). Pearson.
- Wierst, C., Zumbo, B., Rhodes, R. E., Faulkner, G. E., & Beauchamp, M. R. (in press). An

examination of Dweck's psychological needs model in relation to exercise-related well-being. *Journal of Sport and Exercise Psychology*, 1–12. <https://doi.org/10.1123/jsep.2021-0026>

Mplus Code***All Participants Sample, Including Treatment Groups***

VARIABLE: NAMES ARE id cond g1 g2 g3 male age

CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6 CESD12;

MISSING=ALL(999);

USEVARIABLES ARE g1 g2 g3

CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6;

ANALYSIS: ESTIMATOR = MLR;

STITERATIONS=20000;

ITERATION = 40000;

MODEL:

i s q | CESD0@0 CESD1@1 CESD2@2 CESD3@3 CESD4@4 CESD5@5 CESD6@6;

i s q ON g1 g2 g3;

s ON g1 (bs1);

s ON g2 (bs2);

s ON g3 (bs3);

q ON g1 (bq1);

q ON g2 (bq2);

q ON g3 (bq3);

i(v0);

CESD0 - CESD6 (r0-r6);

MODEL CONSTRAINT:

NEW(g1dt1 g1dt2 g1dt3 g1dt4 g1dt5 g1dt6);

$$g1dt1 = (bs1*1 + bq1*1)/\sqrt{(v0+r0/2+r1/2)};$$

$$g1dt2 = (bs1*2 + bq1*4)/\sqrt{(v0+r0/3+r1/3+r2/3)};$$

$$g1dt3 = (bs1*3 + bq1*9)/\sqrt{(v0+r0/4+r1/4+r2/4+r3/4)};$$

$$g1dt4 = (bs1*4 + bq1*16)/\sqrt{(v0+r0/5+r1/5+r2/5+r3/5+r4/5)};$$

$$g1dt5 = (bs1*5 + bq1*25)/\sqrt{(v0+r0/6+r1/6+r2/6+r3/5+r4/6+r5/6)};$$

$$g1dt6 = (bs1*6 + bq1*36)/\sqrt{(v0+r0/7+r1/7+r2/7+r3/7+r4/7+r5/7+r6/7)};$$

NEW(g2dt1 g2dt2 g2dt3 g2dt4 g2dt5 g2dt6);

$$g2dt1 = (bs2*1 + bq2*1)/\sqrt{(v0+r0/2+r1/2)};$$

$$g2dt2 = (bs2*2 + bq2*4)/\sqrt{(v0+r0/3+r1/3+r2/3)};$$

$$g2dt3 = (bs2*3 + bq2*9)/\sqrt{(v0+r0/4+r1/4+r2/4+r3/4)};$$

$$g2dt4 = (bs2*4 + bq2*16)/\sqrt{(v0+r0/5+r1/5+r2/5+r3/5+r4/5)};$$

$$g2dt5 = (bs2*5 + bq2*25)/\sqrt{(v0+r0/6+r1/6+r2/6+r3/5+r4/6+r5/6)};$$

$$g2dt6 = (bs2*6 + bq2*36)/\sqrt{(v0+r0/7+r1/7+r2/7+r3/7+r4/7+r5/7+r6/7)};$$

NEW(g3dt1 g3dt2 g3dt3 g3dt4 g3dt5 g3dt6);

$$g3dt1 = (bs3*1 + bq3*1)/\sqrt{(v0+r0/2+r1/2)};$$

$$g3dt2 = (bs3*2 + bq3*4)/\sqrt{(v0+r0/3+r1/3+r2/3)};$$

$$g3dt3 = (bs3*3 + bq3*9)/\sqrt{(v0+r0/4+r1/4+r2/4+r3/4)};$$

$$g3dt4 = (bs3*4 + bq3*16)/\sqrt{(v0+r0/5+r1/5+r2/5+r3/5+r4/5)};$$

$$g3dt5 = (bs3*5 + bq3*25)/\sqrt{(v0+r0/6+r1/6+r2/6+r3/5+r4/6+r5/6)};$$

$$g3dt6 = (bs3*6 + bq3*36)/\sqrt{(v0+r0/7+r1/7+r2/7+r3/7+r4/7+r5/7+r6/7)};$$

OUTPUT: SAMPSTAT CINTERVAL STANDARDIZED RESIDUAL MODINDICES (3.84);

High Depression Sample, Including Treatment Groups – Free Time Scores

VARIABLE: NAMES ARE id cond g1 g2 g3 male age

CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6 CESD12;

MISSING=ALL(999);

USEVARIABLES ARE g1 g2 g3 CESD0 CESD1 CESD2 CESD3 CESD4 CESD5 CESD6;

ANALYSIS: ESTIMATOR = MLR;

STITERATIONS=5000;

ITERATION = 20000;

MODEL:

i s | CESD0@0 CESD1@1 CESD2* CESD3* CESD4* CESD5* CESD6*;

i s ON g1 g2 g3;

s ON g1 (bs1);

s ON g2 (bs2);

s ON g3 (bs3);

i(v0);

CESD1 - CESD6 (r1-r6);

CESD0@0;

CESD1 WITH CESD2-CESD4;

CESD2 WITH CESD3-CESD5;

CESD3 WITH CESD4-CESD6;

CESD4 WITH CESD5-CESD6;

CESD5 WITH CESD6;

MODEL CONSTRAINT:

NEW(g1dt1 g1dt2 g1dt3 g1dt4 g1dt5 g1dt6);

$g1dt1 = (bs1 * 1) / \sqrt{v0 + r1}$;

$g1dt2 = (bs1 * 2) / \sqrt{v0 + r1/2 + r2/2}$;

$g1dt3 = (bs1 * 3) / \sqrt{v0 + r1/3 + r2/3 + r3/3}$;

$g1dt4 = (bs1 * 4) / \sqrt{v0 + r1/4 + r2/4 + r3/4 + r4/4}$;

$g1dt5 = (bs1 * 5) / \sqrt{v0 + r1/5 + r2/5 + r3/5 + r4/5 + r5/5}$;

$g1dt6 = (bs1 * 6) / \sqrt{v0 + r1/6 + r2/6 + r3/6 + r4/6 + r5/6 + r6/6}$;

NEW(g2dt1 g2dt2 g2dt3 g2dt4 g2dt5 g2dt6);

$g2dt1 = (bs2 * 1) / \sqrt{v0 + r1}$;

$g2dt2 = (bs2 * 2) / \sqrt{v0 + r1/2 + r2/2}$;

$g2dt3 = (bs2 * 3) / \sqrt{v0 + r1/3 + r2/3 + r3/3}$;

$g2dt4 = (bs2 * 4) / \sqrt{v0 + r1/4 + r2/4 + r3/4 + r4/4}$;

$g2dt5 = (bs2 * 5) / \sqrt{v0 + r1/5 + r2/5 + r3/5 + r4/5 + r5/5}$;

$g2dt6 = (bs2 * 6) / \sqrt{v0 + r1/6 + r2/6 + r3/6 + r4/6 + r5/6 + r6/6}$;

NEW(g3dt1 g3dt2 g3dt3 g3dt4 g3dt5 g3dt6);

$g3dt1 = (bs3 * 1) / \sqrt{v0 + r1}$;

$$g3dt2 = (bs3*2)/\sqrt{(v0+r1/2+r2/2)};$$

$$g3dt3 = (bs3*3)/\sqrt{(v0+r1/3+r2/3+r3/3)};$$

$$g3dt4 = (bs3*4)/\sqrt{(v0+r1/4+r2/4+r3/4+r4/4)};$$

$$g3dt5 = (bs3*5)/\sqrt{(v0+r1/5+r2/5+r3/5+r4/5+r5/5)};$$

$$g3dt6 = (bs3*6)/\sqrt{(v0+r1/6+r2/6+r3/6+r4/6+r5/6+r6/6)};$$

OUTPUT: SAMPSTAT CINTERVAL STANDARDIZED RESIDUAL;

Tables*Table S1 - Missing Surveys at Item and Survey Levels.*

Week	Non-missing Survey (All items complete)	Non-Completely Missing Survey (Missing ≥ 1 items)	Completely Missing Surveys
0	327	7	0
1	306	9	19
2	299	5	30
3	290	2	42
4	287	6	41
5	276	6	52
6	284	6	44

Table S2 - Fit Indices for the Primary SEM Models.

Measure	All Participants - Quadratic Time	High Depression – Free Time Scores
Root Mean Square Error of Approximation, Est. (.90CI)	0.049 (0.027, 0.069)	0.042 (0.000, 0.081)
CFI	0.983	0.991
Standardized Root Mean Square Residual	0.026	0.044

Table S3 - Income Grouping.

Income Values	Recoded
1 - 10,000 10,001 - 20,000 20,001 - 30,000 30,001 - 40,000	0 – 40,000
40,001 - 50,000, 50,001 - 60,000, 60,001 - 70,000, 70,001 - 80,000	40,001 – 80,000
80,001 - 90,000, 90,001 - 100,000, 100,001 - 110,000, 110,001 - 120,000	80,001 – 120,000
120,001 - 130,000, 130,001 - 140,000, 140,001 - 150,000, 150,001 - 160,000	120,001 – 160,000
160,001 - 170,000, 170,001 - 180,000, 180,001 - 190,000, 190,001 - 200,000, 200,001 - 210,000, 210,001 - 220,000,	160,000 +

220,001 - 230,000, 230,001 - 240,000, 240,001 - 250,000, 250,001+	
Do not know / prefer not to answer, [Missing]	No response

Table S4 - Employment Grouping.

Current Employment	Other (Text Response)	Coded Value
Other	Full time work with one furlough day due to covid	Full Time
Other	teacher in spring semester	Full Time
Other	Teacher summer vacation	Full Time
Working full-time		Full Time
Working full-time,Other	on summer vacation	Full Time
Working full-time,Self-employed		Full Time
Working full-time,Self-employed ,Other	I'm working full time at a company, but also self-employed teaching (Not full-time-self-employed)	Full Time
Homemaker		Not Working
Looking for work; unemployed		Not Working
Looking for work; unemployed,Homemaker		Not Working
Looking for work; unemployed,Other	housewife	Not Working
Looking for work; unemployed,Other	working - unpaid	Not Working
Looking for work; unemployed,Temporarily laid off		Not Working
Maternity or sick leave (volunteered)		Not Working
Other	Leave of absence	Not Working

Other	Self employed but no work available	Not Working
Other	Stay at home parent	Not Working
Self-employed ,Looking for work; unemployed,Permanently disabled (volunteered),Other	Home based business not in market due to covid	Not Working
Self-employed ,Temporarily laid off		Not Working
Temporarily laid off		Not Working
Temporarily laid off,Maternity or sick leave (volunteered)		Not Working
[Missing]		Other
Don't know/not sure		Other
Other	Will be laid off during this study	Other
Other		Other
Prefer not to say		Other
Self-employed		Other
Working part-time		Part Time
Working part-time,Homemaker		Part Time
Working part-time,Other	on CEWS	Part Time
Working part-time,Self-employed		Part Time
Working part-time,Self-employed ,Homemaker		Part Time
Full-time student		Student

Looking for work; unemployed,Full-time student		Student
Looking for work; unemployed,Part-time student		Student
Looking for work; unemployed,Temporarily laid off,Part-time student		Student
Part-time student		Student
Self-employed ,Full-time student		Student
Temporarily laid off,Full-time student		Student
Temporarily laid off,Part-time student		Student
Working full-time,Full-time student		Student
Working full-time,Part-time student		Student
Working part-time,Full-time student		Student
Working part-time,Full-time student,Other	Support from parents	Student
Working part-time,Part-time student		Student

Table S5 - Education Grouping.

Educational Status	Recoded
High school diploma or a high school equivalency certificate	High School or less
College, CEGEP or other non-university certificate or diploma (other than trades certificates or diplomas),	College, trade school or certificate
Trade certificate or diploma,	College, trade school or certificate
University certificate or diploma below the bachelor's level	College, trade school or certificate
Bachelor's degree (e.g. B.A., B.Sc., LL.B.)	Bachelor or equivalent
University certificate, diploma, degree above the bachelor's level	Postgraduate / professional training
Prefer not to answer,	No response
[Missing]	No response

Note: "Less than high school diploma or its equivalent" was an option, but no participant selected it.

Table S6 - Cultural Background Grouping.

Cultural Background	Other (Text Response)	Coded Value
Chinese		Asian
Chinese,Filipino		Asian
Chinese,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Asian
Filipino		Asian
Filipino,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Asian
Japanese		Asian
Korean		Asian
Other - please specify	Taiwanese	Asian
South Asian (e.g., East Indian, Pakistani, Sri Lankan, etc)		Asian
South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Asian
Aboriginal decent (e.g., North American Indian, Métis or Inuit (Eskimo))		Other
Aboriginal decent (e.g., North American Indian, Métis or Inuit (Eskimo)),Black (e.g., African, Haitian, Jamaican, Somali, etc)		Other
Aboriginal decent (e.g., North American Indian, Métis or Inuit (Eskimo)),White		Other

Arab		Other
Black (e.g., African, Haitian, Jamaican, Somali, etc)		Other
Latin American		Other
Other - please specify		Other
Other - please specify	european / mixed	Other
Other - please specify	Doesn't let you choose more than one. Aboriginal and European.	Other
Other - please specify	Afghan	Other
Other - please specify	Austrian	Other
South Asian (e.g., East Indian, Pakistani, Sri Lankan, etc),Latin American		Other
West Asian (e.g., Iranian, Afghan, etc)		Other
White,Arab		Other
White,Chinese		Other
White,Chinese,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Other
White,Filipino		Other
White,Japanese		Other
White,Latin American		Other
White,Other - please specify	Jewish	Other

White,Other - please specify	European	Other
White,Other - please specify	Ashkenazi Jewish	Other
White,Other - please specify	Scandinavian	Other
White,Other - please specify	Dutch	Other
White,South East Asian (e.g., Vietnamese, Cambodian, Malaysian, Laotian, etc)		Other
Other - please specify	Italian	White
Other - please specify	IRISH	White
White		White

Note: As cultural backgrounds were asked as check boxes, many participant selected multiple options, making it difficult to define groups. Therefore, participants were split into White, Asian and Other, as White and Asian participants made up a large majority of the sample.

Table S7 - Marital Status Groupings.

Marital Status	Recoded
Married	Married
Living common-law	Married
Separated	No Longer Married
Divorced	No Longer Married
Widowed	No Longer Married
Single, never married	Single
Prefer not to answer	Other
[Missing]	Other

Table S8 - Weekly Survey Response Rate, N (%), by Experimental Grouping.

	Week						
	0	1	2	3	4	5	6
WLC	83 (100%)	80 (96%)	77 (93%)	80 (96%)	76 (92%)	77 (93%)	74 (89%)
HIIT	82 (100%)	77 (94%)	76 (93%)	69 (84%)	72 (88%)	64 (78%)	68 (83%)
Yoga	86 (100%)	83 (97%)	76 (88%)	74 (86%)	76 (88%)	75 (87%)	67(78%)
HIIT+Yoga	83 (100%)	75 (90%)	75 (90%)	69 (83%)	69 (83%)	66 (80%)	84 (82%)
	Week						
	0	1	2	3	4	5	6
WLC	83 (100%)	80 (96%)	77 (93%)	80 (96%)	76 (92%)	77 (93%)	74 (89%)
HIIT	82 (100%)	77 (94%)	76 (93%)	69 (84%)	72 (88%)	64 (78%)	68 (83%)
Yoga	86 (100%)	83 (97%)	76 (88%)	74 (86%)	76 (88%)	75 (87%)	67(78%)
HIIT+Yoga	83 (100%)	75 (90%)	75 (90%)	69 (83%)	69 (83%)	66 (80%)	84 (82%)

Table S9 – Average Number of Minutes of DownDog Exercises Completed by Week and Experimental Group

Week	HIIT	Yoga	Combo
1	63.95 (29.73)	73.13 (43.43)	80.72 (48.48)
2	64.44 (35.03)	78.43 (46.70)	86.04 (53.80)
3	59.39 (35.49)	76.76 (53.98)	78.31 (45.19)
4	58.90 (36.08)	73.44 (59.71)	74.20 (55.49)
5	53.15 (36.85)	68.57 (50.81)	72.86 (56.29)
6	46.11 (39.02)	68.60 (57.62)	70.97 (62.29)

Table S10 – Weekly Average (SD) Minutes of Moderate-to-Vigorous Physical Activity (Not Including App Usage) for Each Group

Group	Week	Moderate	Vigorous	MVPA
Control	0	16.73 (33.25)	0.00 (0.00)	18.51 (34.53)
Control	1	40.71 (81)	7.49 (21.95)	49.04 (90.77)
Control	2	42.68 (63.32)	9.32 (29.21)	52.57 (72.50)
Control	3	44.32 (60.7)	4.34 (15.37)	47.15 (65.95)
Control	4	35.42 (48.19)	7.47 (21.26)	43.38 (57.89)
Control	5	44.17 (67.25)	6.1 (18.66)	49.91 (72.59)
Control	6	71.67 (83.48)	6.16 (16.89)	74.52 (81.26)
HIIT	0	19.72 (34.26)	0.30 (2.46)	19.06 (34.79)
HIIT	1	40.00 (52.03)	6.38 (19.11)	48.49 (60.39)
HIIT	2	45.36 (59.22)	15.40 (35.00)	61.88 (74.94)
HIIT	3	45.17 (66.58)	12.71 (26.82)	59.12 (79.29)
HIIT	4	59.25 (82.2)	14.32 (31.79)	71.29 (89.25)
HIIT	5	41.86 (55.51)	15.26 (31.00)	57.02 (73.43)
HIIT	6	49.33 (80.68)	13.71 (29.41)	59.69 (88.17)
Yoga	0	26.86 (38.26)	1.01 (4.25)	24.57 (36.49)
Yoga	1	56.88 (89.41)	11.46 (31.36)	68.66 (107.56)
Yoga	2	44.42 (58.8)	7.17 (21.77)	52.58 (67.24)
Yoga	3	47.84 (60.56)	14.18 (25.82)	58.15 (69.24)
Yoga	4	56.94 (68.03)	16.54 (33.89)	73.38 (81.59)
Yoga	5	73.46 (106.01)	11.42 (24.41)	85.07 (116.04)

Yoga	6	51.58 (72.44)	15.91 (30.62)	66.75 (92.63)
Combo	0	18.87 (35.98)	0.99 (4.11)	20.46 (37.72)
Combo	1	46.15 (97.01)	10.73 (29.82)	58.07 (103.51)
Combo	2	32.46 (45.73)	10.76 (27.46)	43.43 (54.73)
Combo	3	56.52 (70.15)	13.52 (25.7)	67.59 (74.84)
Combo	4	49.41 (68.91)	17.70 (33.67)	65.78 (85.26)
Combo	5	66.25 (99.03)	10.79 (26.15)	75.20 (112.14)
Combo	6	53.49 (74.67)	12.66 (29.28)	59.75 (78.46)

Table S11 - Estimates for Trajectories for HIIT (A) and Comparisons with WLC, Yoga and HIIT+Yoga Groups (B1-3).

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I; estimated baseline), slope (S; time), and quadratic (Q; time²) terms for HIIT</u>			
I	10.60	0.64	9.34, 11.86
S	-0.93	0.27	-1.45, -0.40
Q	0.10	0.04	0.03, 0.18
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
WLC vs HIIT	-0.26	0.87	-1.97, 1.46
Yoga vs HIIT	-0.59	0.89	-2.34, 1.16
HIIT+Yoga vs HIIT	-1.53	0.87	-3.23, 0.17
<u>B2. Differences in estimates for S</u>			
WLC vs HIIT	0.77	0.38	0.04, 1.51
Yoga vs HIIT	0.07	0.40	-0.72, 0.86
HIIT+Yoga vs HIIT	-0.14	0.40	-0.93, 0.65
<u>B3. Differences in estimates for Q</u>			
WLC vs HIIT	-0.11	0.06	-0.22, 0.00
Yoga vs HIIT	-0.03	0.06	-0.15, 0.08
HIIT+Yoga vs HIIT	0.01	0.06	-0.11, 0.12

Note: Results from the SEM. model estimating intercept, slope, and quadratic term for HIIT group (Section A) and comparisons of these estimates with those of the three other groups (WLC, Yoga, HIIT+Yoga; Section B). Bold text denotes $p < .05$

Table S12 - Estimates for Trajectories for HIIT+Yoga (A) and Comparisons with WLC, HIIT and Yoga Groups (B1-3).

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I; estimated baseline), slope (S; time), and quadratic (Q; time²) terms for HIIT+Yoga</u>			
I	9.07	0.58	7.93, 10.20
S	-1.07	0.30	-1.65, -0.48
Q	0.11	0.05	0.02, 0.19
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
WLC vs HIIT+Yoga	1.27	0.83	-0.35, 2.89
HIIT vs HIIT+Yoga	1.53	0.87	-0.17, 3.23
Yoga vs HIIT+Yoga	0.94	0.85	-0.72, 2.59
<u>B2. Differences in estimates for S</u>			
WLC vs HIIT+Yoga	0.92	0.39	0.14, 1.69
HIIT vs HIIT+Yoga	0.14	0.40	-0.65, 0.93
Yoga vs HIIT+Yoga	0.21	0.42	-0.62, 1.04
<u>B3. Differences in estimates for Q</u>			
WLC vs HIIT+Yoga	-0.11	0.06	-0.23, 0.01
HIIT vs HIIT+Yoga	-0.01	0.06	-0.12, 0.11
Yoga vs HIIT+Yoga	-0.04	0.06	-0.16, 0.09

Note: Results from the SEM model estimating intercept, slope, and quadratic term for HIIT+Yoga group (Section A) and comparisons of these estimates with those of the three other groups (WLC, HIIT, Yoga; Section B). Bold text denotes $p < .05$

Table S13 - Estimates for Trajectories for Yoga (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1-3).

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I; estimated baseline), slope (S; time), and quadratic (Q; time²) terms for Yoga</u>			
I	10.01	0.62	8.78, 11.23
S	-0.86	0.31	-1.45, -0.26
Q	0.07	0.05	-0.02, 0.16
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
WLC vs Yoga	0.34	0.86	-1.34, 2.02
HIIT vs Yoga	0.59	0.89	-1.16, 2.34
HIIT+Yoga vs Yoga	-0.94	0.85	-2.59, 0.72
<u>B2. Differences in estimates for S</u>			
WLC vs Yoga	0.70	0.40	-0.08, 1.49
HIIT vs Yoga	-0.07	0.40	-0.86, 0.72
HIIT+Yoga vs Yoga	-0.21	0.422	-1.04, 0.62
<u>B3. Differences in estimates for Q</u>			
WLC vs Yoga	-0.08	0.06	-0.20, 0.05
HIIT vs Yoga	0.03	0.06	-0.08, 0.15
HIIT+Yoga vs Yoga	0.04	0.06	-0.09, 0.16

Note: Results from the SEM. model estimating intercept, slope, and quadratic term for HIIT group (Section A) and comparisons of these estimates with those of the three other groups (WLC, HIIT, HIIT+Yoga; Section B). Bold text denotes $p < .05$

Table S14 - Effect Sizes for Model with all Individuals and Model including only those with High Depressive Symptoms at Baseline.

All Participants						
Time	HIIT		Yoga		HIIT+Yoga	
	Effect Size	95% CI	Effect Size	95% CI	Effect Size	95% CI
Week 1	-0.12	[-0.23, -0.01]	-0.11	[-0.23, 0.01]	-0.14	[-0.26, -0.03]
Week 2	-0.20	[-0.39, -0.01]	-0.20	[-0.39, 0.00]	-0.24	[-0.44, -0.05]
Week 3	-0.24	[-0.48, 0.00]	-0.25	[-0.49, -0.02]	-0.31	[-0.54, -0.07]
Week 4	-0.24	[-0.50, 0.02]	-0.28	[-0.53, -0.04]	-0.33	[-0.58, -0.08]
Week 5	-0.21	[-0.48, 0.06]	-0.29	[-0.52, -0.06]	-0.31	[-0.55, -0.07]
Week 6	-0.14	[-0.44, 0.16]	-0.27	[-0.51, -0.04]	-0.25	[-0.51, 0.00]
Subpopulation with High Depressive Symptoms						
Time	HIIT		Yoga		HIIT+Yoga	
	Effect Size	95% CI	Effect Size	95% CI	Effect Size	95% CI
Week 1	-0.41	[-0.69, -0.14]	-0.40	[-0.67, -0.14]	-0.44	[-0.71, -0.18]
Week 2	-0.83	[-1.38, -0.28]	-0.80	[-1.33, -0.28]	-0.89	[-1.43, -0.35]

Week 3	-1.20	[-2.01, -0.40]	-1.17	[-1.94, -0.39]	-1.29	[-2.07, -0.51]
Week 4	-1.57	[-2.62, -0.52]	-1.52	[-2.53, -0.51]	-1.68	[-2.70, -0.67]
Week 5	-1.94	[-3.21, -0.66]	-1.88	[-3.11, -0.65]	-2.08	[-3.32, -0.84]
Week 6	-2.34	[-3.87, -0.81]	-2.27	[-3.74, -0.80]	-2.51	[-4.00, -1.01]

Note: All effect sizes are compared to the WLC group.

Table S15 - Estimates for Trajectories for WLC (A) and Comparisons with WLC, HIIT and HIIT+Yoga Groups (B1 & B2) in Participants with High Levels of Depression Symptoms at Baseline.

	Estimate	SE	95% CI
<u>A. Estimates, SE, and 95% CI for intercept (I) and slope (S) terms for WLC</u>			
I	14.41	0.56	13.32, 15.51
S	-1.18	0.44	-2.05, -0.31
<u>B. Estimates for differences between each group and WLC</u>			
<u>B1. Differences in estimates for I</u>			
HIIT vs WLC	1.04	0.85	-0.62, 2.70
Yoga vs WLC	0.52	0.81	-1.06, 2.10
HIIT+Yoga vs WLC	-0.26	0.83	-1.89, 1.38
<u>B2. Differences in estimates for S</u>			
HIIT vs WLC	-2.06	0.68	-3.39, -0.73
Yoga vs WLC	-2.00	0.65	-3.28, -0.72
HIIT+Yoga vs WLC	-2.21	0.67	-3.52, -0.90

Note: Results from the SEM model, including only those with high levels of depressive symptoms at baseline, estimating intercept and slope for waitlist control (Section A) and comparisons of these estimates with those of the three active groups (HIIT, Yoga, HIIT+Yoga; Section B). Bold text denotes $p < .05$

Figures

Figure S1 - SEM Path Diagram for Model including all Participants.




Figure S2 - SEM Path Diagram for Model including Participants with High Baseline Depressive Symptoms.

